arXiv:cond-mat/0601210v1 [cond-mat.stat-mech] 10 Jan 2006

Viscosity from Newton to Nonequilibrium Statistical Mechanics[*]

Sébastien Viscardy[†]
Center for Nonlinear Phenomena and Complex Systems,
Université Libre de Bruxelles,
Campus Plaine, Code Postal 231, B-1050 Brussels, Belgium

Abstract

This paper is devoted to the transport process of momentum, the viscosity. We outline the historical development of hydrodynamics, the branch of physics in which the shear and bulk viscosities are expressed. We show the transition between the mechanical philosophy claiming that the conservation of energy is coupled to cyclic, reversible dynamics, and the birth of a new science, thermodynamics, in which the conservation of energy (the first law) is associated with the irreversibility (the second law). Through the 19th century, the kinetic theory of gases was developed and accumulated successes establishing relationships between the microscopic dynamics of the atoms and molecules, and macroscopic properties, among which the viscosity. In this context appeared the paradox of irreversibility to which Boltzmann was the first to provide an explanation based on a *statistical* interpretation of the entropy. We finally outline the history of the theory of chaos and show how the chaotic character of the microscopic dynamics provides a *dynamical* explanation of the irreversible processes such as transport, and in particular, the viscosity. In this context, we present three successfull approaches developed the last two decades: the *thermostated-system method*, the *escape-rate formalism* and the *hydrodynamic-mode method*.

---

[*] This paper consists in a revision of the introduction of the PhD thesis presented on the 21th of September, 2005, entitled : *Viscosity and Microscopic Chaos : The Helfand-moment Approach* (Viscardy, 2005).

[†] e-mail: sviscard@ulb.ac.be

1 Introduction

The nineteenth century was rich in scientific discoveries and in new concepts. In particular, as regards the subject discussed in this paper, the nineteenth century appears as a turning point. Indeed, the industrial revolution gave rise to the development of a new science, *thermodynamics*, which emphasized the *irreversible* character of macroscopic phenomena (such as heat dissipation). On the other hand, the development of chemistry and, later, of the kinetic theory of gases, transformed the concept of matter from a continuous to a discrete description. In other words, matter should be considered as made of very small particles: the atoms and the molecules. However, the motion of these entities is governed by the *reversible* equations of Newton. As a consequence appeared the famous paradox opposing macroscopic and microscopic descriptions, which has given rise after the seminal works of Maxwell and Boltzmann to a large number of contributions addressing this paradox.

Maxwell was the first to introduce viscosity in the framework of kinetic theory and nonequilibrium statistical physics. Since then, it has remained a driving concept in these fields. We therefore devote the paper to the case of viscosity as a transport property.

In the sixties a new revolutionary field of the science of the twentieth century was born : Chaos. Sometimes compared to the scientific revolutions implied by quantum mechanics and the relativity, the phenomenon of chaos has been discovered in most of the natural sciences such as physics, chemistry, meteorology and geophysics. Our intent is to show the role of microscopic chaos in statistical mechanics of irreversible processes.

The paper is organized as follows: we will start in section 2 with the presentation of the discovery of the concept of irreversibility as a fundamental property in nature. The development of hydrodynamics as the domain of physics being particularly concerned by the process of viscosity will be considered in section 3. Further, in section 4, we focus on the kinetic theory that has a long history and has played an important role in the establishment of the statistical mechanics, that is to say, in the understanding of the relation between the microscopic and macroscopic behaviors. Finally, in section 5, we cover the brief history of the development of the theory of chaos and of dynamical systems which provides a better understanding of the origin of the irreversible processes.

2 Irreversibility

The history of modern science holds its richness in the variety of conceptions which were developed successively throughout the years[1]. In the 17th century, it was the doctrine of the *clockmaker God* or *mechanical philosophy* which prevailed above the others.

The development of these ideas began with the work of Boyle (1627-1691) and others. According to his doctrine, Boyle stated that the world works like a clock: once created and energized, it can run forever in a deterministic way and without any need for a divine intervention. To ensure that this “clockwork universe” never runs down, Descartes (1596-1650) more or less intuitively introduced the statement that the total amount of motion in the world must remain constant. He defined this quantity as the *scalar* momentum mv (quantity of matter multiplied by its scalar speed[2]) (Blackwell, 1966). However, as Descartes himself later observed, experiments did not confirm his enunciation of the conservation law of motion. Huygens (1629-1695) corrected it in 1668 by modifying it into the *vector* form. Huygens worked also on the problem of collisions. This revised law allowed him to claim that the vector sum of the product of the mass by the vector velocity $m\mathbf{v}$ remains unchanged after a collision even if it was inelastic and with dissipation of energy after the collision itself. Introducing this modification, it then appears that the world could stop after a certain amount of time. This was totally contradictory with the clockworld concept.

The only way to avoid this possibility was to postulate that the matter is composed of elastic particles. If the macroscopic objects lose some motion after a collision, it is only in appearance because the motion is transferred to the invisible particles of the objects. Therefore, the idea emerged that heat is related to a rapid motion of the invisible parts inside the macroscopic bodies.

In the 18th century, the study of the motion of bodies and its change after collisions was of great importance. Moreover, in addition to this study, another quantity so-called *vis viva*, mv^2, was introduced for the first time. Besides, its conservation law in elastic collisions was remarkably proved by the same man who helped formulate the conservation of momentum: Huygens. This major contribution induced the first step to the central quantity acknowledged in mechanics: the energy. The concept of *vis viva* was used up to the 19th century, when the factor $\frac{1}{2}$ was added and it then became "kinetic energy".

---

[1] For general informations on this section, see for example Holton and Brush (1985) and Brush (1983) .

[2] More precisely, Descartes considered the volume of the bodies instead of their mass, the volume being for him the real measure of quantity of matter to be considered, as pointed out by Blackwell (1966) .

The world-machine concept is often attributed to Newton (1642-1727) because of the importance played by his brilliant *Principia* published in 1687 (Newton, 1999). In his masterpiece, the Newtonian dynamics is depicted for the first time although it clearly appeared that the author was really opposed to it as he pointed out in his *Opticks* (Newton, 1952) published in 1704

> By reason of the [...] weakness of elasticity in solids, motion is much more apt to be lost than got, and is always upon the decay.

Therefore, in a certain sense, he suggested the existence of the dissipation of motion in the world. However, it should be specified that this viewpoint has to be considered in the light of the polemic between Newton[3] and Leibniz concerning the role of God (Koyré, 1957).

During the 18th century the concept of the "Newtonian world-machine" continued to dominate the thought of scientists. This century is also the one of the birth of Geology. It began with the question of the temperature of Earth's interior (Brush, 1994). Yet in 1693 Leibniz thought that Earth was originally hotter and cooled down, at least on the outside. Later Buffon (1707-1788) studied this question by leading some experiments on the cooling of heated spheres of iron. By considering a molten Earth at the origin he hence found that our planet was about 75,000 years (de Buffon, 1774). At the opposite, the founder of Geology James Hutton (1726-1797) disagreed with this theory of cooling. Defending his "Uniformitarism" (principle telling that the geological processes in the past have to be explained by using only the laws and physical processes that can now be observed) and accepting the hypothesis that the interior of the Earth is much hotter than its surface, he thought this situation had been like that forever. For him the geological processes are cyclic: alternance of periods of erosion and denudation implying the destruction of the mountains, and periods of uplifts of new continents (thanks to the subterranean fires). Actually one of his disciples Playfair promoted the uniformitarist viewpoint by citing the mathematical works by Lagrange and Laplace showing the cyclic movement of the planets around the sun. This position was perpetuated by Lyell (1797-1875) during the 19th century.

In the early 19th century, the first mathematical theory describing the propagation of heat was developed. Indeed, the French mathematician

---

[3] More precisely, his student and faithful friend Clarke (1675-1729).

Fourier (1768-1830) was interested by the problem of the cooling of the Earth. This was the main motivation that lead him to study the heat conduction in solids. In 1819, in his *Mémoire sur le refroidissement séculaire du globe terrestre* (Fourier, 1819), he came up with an equation which presented a significant feature: unlike Newton's laws of motion, Fourier's equation is *irreversible*. Any system in which a temperature difference exists presents a heat flow from the high to low temperature. This was the discovery of an equation describing processes in which the past does not play the same role as the future. His theory therefore presents an important turning point in the history of physics due to the powerful analysis he developed (what we now call Fourier's analysis), and because it is explicitly based on a postulate of irreversibility. In the first half of the 19th century the eventual contradiction between the reversible Newtonian dynamics and the irreversible heat conduction did not appear because the most studied problem at this time was at the phenomenological level and the Newtonian mechanics had already been successfully applied to problems with dissipative forces (friction, etc.). The contradiction appeared during the first attempt to explain the macroscopic processes in terms of the assumed reversible Newtonian dynamics of the particles composing the system.

A few years after Fourier's work on the propagation of heat, Carnot (1796-1832) published in 1824 his essay *Réflexions sur la force motrice du feu* (Carnot, 1824). Driven by engineering ideas, he focused his research on the question of the limited efficiency of the steam engine. By using the *caloric theory*[4] appearing during the 18th century and developed by Lavoisier (1743-1794), Carnot concluded that a difference of temperature existing between two bodies gives the possibility of doing work by allowing heat to expand a gas as the heat flows from the hot body to the cold one[5]. However, an engine wrongly designed results in loosing the chance of doing work that might have been done. Therefore, proving that heat always flows from hot to cold bodies in the engine as it does anywhere in nature, and discovering that this equalization represents a loss of the opportunity to produce mechanical work, Carnot pointed out the existence of *dissipation.* More important, the discovery made first by Carnot in 1831, then Mayer (1814-1878) in 1842 (Mayer, 1842) and later by Joule (1818-1889) (Joule, 1847), brought up a fundamental statement: work and heat are actually two different expressions of the same quantity: the energy. Indeed, although heat and mechanical work seem to be two independent quantities, we now understand that heat can be transformed into work and vice versa. In other terms, the energy of an isolated system remains constant. A science of transformations was born: *thermodynamics.* This equivalence between heat Q and work W consists in the so-called *first law of thermodynamics*: the conservation of the internal energy E of a closed system: $dE = dQ + dW$.

---

[4] The caloric theory supposed that heat is a fluid composed of particles independent of the rest of matter. These particles repel each other but are attracted to the particles of ordinary matter. According to this theory it has to be an eventual material substance and thus should be limited. Hence the increase of the temperature of water by rotating a drill should be due to the transfer of the so-called *caloric* from the drill to the water. But, in 1798, Thompson (1753-1814) realized that one can produce heat without any limit. He then concluded that heat is not a chemical substance or a material substance but is the expression of a movement. However, since he did not propose any alternative theory of heat, the caloric theory had an important influence until the 1830's and the works done in particular by Joule.

[5] Actually later, Carnot abandoned the caloric theory.

As we have seen, the original problem giving birth to thermodynamics is the separation between the concepts of conservation and reversibility. In mechanical transformations, the conservation of energy (at the beginning the *vis viva*) coincides with the idea of reversibility. On the other hand, the physico-chemical transformations can preserve the energy while they cannot be reversed. Hence it became necessary to define a quantity characterizing this irreversibility. In 1854 Clausius (1822-1888) introduced the quantity dQ/T as a measure of the quantity of work lost during the transfer of heat from a hot to a cold body (Clausius, 1854; Clausius, 1856). Later, in 1865, he gave it the famous name *entropy*[6] (Clausius, 1865). Thanks to this new quantity, the *second law of thermodynamics* can now be stated very simply: the entropy of an isolated system always tends to increase. We thus have a well-established principle regarding the irreversible processes. We may therefore state that the two laws of thermodynamics combine the two concepts of conservation of energy and of nonreversibility of macroscopic phenomena.

As we will later see in section 4, the atomic theory increased its influence during the 19th century. The development of the kinetic theory of gases played an important role to prove the discrete character of matter. One of the most important leaders in this area was Boltzmann (1844-1906). He particularly proposed a more general definition of entropy, in terms of the probabilities of molecular arrangements, that can change even when there is no heat flow (e.g. the system become more disordered). His theory based on a statistical description was a first understanding of the link between the atomic level and the macroscopic phenomena, that is between the reversible microscopic dynamics and the macroscopic irreversible processes.

---

[6] The ethymology of this word expressed in the most clearly way the essence of thermodynamics: indeed it comes from the Greek words $\epsilon\nu\epsilon\rho\gamma\epsilon\iota\alpha$ (energy) and $\tau\rho o\pi\eta$ (transformation).

In the nineteenth-century society, in which social and economical activities, science and technology induced a progress never observed before in the history of Humanity, arose the idea of the evolution of species with Lamark (1744-1829) and later with Darwin (1809-1882). According to Darwin the species that we know today are the result of a long process of natural selection. This mechanism has induced the production of increasingly complex and organized beings, from the unicellular bacteries to the mammals such as the human beings. Whereas thermodynamics introduced a quantity measuring the continuous growth of disorder, biology put in evidence the continuous growth of order, of organization in the biological world. As a result, these two points of view influenced some philosophers: on the one hand, Adams (1838-1918), influenced by the discovery of constant dissipation of energy in the universe, expressed a pessimistic vision concerning the future of the society, as argued in *The Degradation of the Democratic Dogma* (Adams, 2004), and made the second law of thermodynamics an explicit basis for the tendency of history (Burich, 1987); on the other hand, the Darwinism strongly influenced philosophers like Spencer (1819-1903) who thought that all in universe goes gradually from a state of confused homogeneity to a differentiated heterogeneous state (Freeman, 1974; Rousseau, 1945). These two points seemed to be contradictory. One had yet to wait for the development of the nonequilibrium thermodynamics, especially attributed to Prigogine (1917-2003) and his coworkers, to resolve this apparent contradiction. Indeed they showed that, out of thermodynamic equilibrium, i.e. in open systems, matter is able to exhibit self-organization, although the internal production of entropy is absorbed by the environment (Glansdorff and Prigogine, 1971; Nicolis and Prigogine, 1977). These recent works have pointed out the importance of nonequilibrium phenomena and have led to important advances on the properties of nonequilibrium systems. It is in this general context that this thesis is written.

3 Hydrodynamics and viscosity

The vital importance of water throughout all civilizations induced a special interest for the study of the properties and behaviors of fluids (in particular of water). The earliest quantitative application of “real fluid” or viscosity effects was by the ancient Egyptian Amenemhet (∼1600 BC). He made a 7 degree correction to the drainage angle of a water clock to account for the viscosity change of water with temperature (which can be significant between day and night in this region). And in general the early centres of civilization in Egypt, Mesopotamia, India and China systematically used various machines for irrigation and water supply (Blair, 1944). In particular, Archimedes (287-212 BC), is not only considered as the father of hydrostatics for his famous law, but has also developed the so-called Archimedes screw, a water elevating machine, which has been used for different purposes (e.g. to extract water from rivers). Notice also that Heron of Alexandria ($\sim$ 10-75) treated in *Mechanica* the problem of friction between a body and a plane and, considering a horizontal, frictionless plane, he said

> We demonstrate that a weight in this situation can be moved by a force less than any given force[7].

The earliest extensive treatise devoted to practical hydraulics is a manual on urban water supply written by Roman soldier and engineer Sextus Julius Frontius (first century BC), inspector of the aqueducts and the public fontains in Rome. He noted that the amount of water discharged through an orifice in a given interval of time depends, not only on the size of the orifice, but also on its depth h below the surface of the water in the reservoir from which the water is supplied. Starting upon the results of Galileo's experiments with falling bodies, Torricelli (1608-1647) came to the conclusion that the velocity v of the water exiting out of the reservoir is proportional to the square root of the h. This is *Torricelli's theorem* which has been mathematically expressed as $v = \sqrt{2gh}$ later by Daniel Bernoulli (1700-1782). On the other hand, the centuries since the *Renaissance* have been characterized by the major influence of the engineerings on the development of physics. More specifically, the construction of bridges and canals induced a lot of theoretical studies on the flow of fluids. By observing the behavior of the flow of water in rivers, da Vinci (1425-1519) already came to the conclusion that, when the river becomes shallower or narrower, the water flows faster. Later Castelli (1577-1644) confirmed this result and gave the so-called *velocity-area law*: $vA =$ const, where v is the velocity of water and A the cross-sectional area of the flow. It consists of the first idea of continuity of flow (for an incompressible fluid), which was later developed by Euler.

The term *hydrodynamics* was first used by Daniel Bernoulli as the title of his book *hydrodynamica* published in 1738 (Bernoulli, 1738). His theory was original because he was the first scientist to combine the sciences of hydrostatics (considering the pressure) and hydraulics (considering the motion of fluids). From the conservation of *vis viva* he reached the famous so-called *Bernoulli's principle* relating the velocity of flow at a point in a pipe to the pressure there: $\frac{\rho v^2}{2} + P = \text{const}$. But in the modern sense, hydrodynamics began with the work of d'Alembert (1717-1783) and especially Euler (1707-1783).

---

[7] Cited by Russo (2004).

During his Berlin period as an engineer, Euler was in charge of the construction of canals, the supply of water for King's Sans-Soucis Palace, and the improvement the water turbine. However, his strong interest for mathematics, mechanics and physics was much greater. Hydrodynamics was among his numerous topics of interest. In 1755, he derived the fundamental equations of hydrodynamics (Euler, 1755) by introducing the new and main concept of *fluid particle*[8]. First he proposed the modern and general form for the equation of continuity (1), which expresses the conservation of matter[9]

$$\frac{d\rho}{dt} + \rho \frac{\partial v_j}{\partial r_j} = 0 \; . \tag{1}$$

where ρ is the mass density and v_i the i-component of the velocity of the considered fluid particle. Furthermore, considering a small parallelepiped and pressure P acting on its different faces, by starting from the Newton's second law, he derived the general equations for the motion of an ideal fluid, the so-called *Euler's equations of motion*

$$\rho \frac{dv_i}{dt} = \rho \left(\frac{\partial v_i}{\partial t} + v_j \frac{\partial v_i}{\partial r_j} \right) = -\frac{\partial P}{\partial r_i} + f_i \; , \tag{2}$$

where f_i is the i-component of the external force (gravity,...).

But these equations do not present the property of fluids which was more and more of interest at the beginning of the nineteenth century, that is the internal friction, or the *viscosity*[10] in modern terminology. Indeed, the dissipative feature of the fluid dynamics does not appear in Euler's equations and it was not until then considered as a fundamental property in nature.

---

[8] A fluid particle is imagined as an infinitesimal body, small enough to be treated mathematically as a point, but large enough to possess such physical properties as volume, mass, density, and so on.

[9] We use the Einstein's convention of summation over repeated indices: $v_j \frac{\partial v_i}{\partial r_j}$ has to be understood as $\sum_j v_j \frac{\partial v_i}{\partial r_j}$.

[10] The word viscosity derives from the latin word *viscum* for mistletoe. The mistletoe berries produce a viscous glue which was used to lime birds. Viscosity is expressed in $Pa.s$ or in *Poise* ($10^{-1} Pa.s$) in the honor of Poiseuille (1797-1869).

The first studies on this property go back already to Newton's Book II of his *Principia* (Newton, 1999). His theory stated that, if a portion of fluid is kept moving, the motion gradually communicates itself to the rest of the fluid. He ascribed to this effect the term *defectus lubricitatis*, that is, a *lack of slipperiness*. It is nothing but viscosity. Considering two particles of a viscous flow sliding one over another, then there exists friction, or viscous resistance along the surface of contact. The force of this resistance per unit area is nowadays known as the *shear stress* τ. The hypothesis proposed by Newton[11] is that the shear stress depends on the speed with which the two layers slide one over another. By extension, we obtain that the shear stress is proportional to the velocity gradient in the perpendicular direction of the movement of the fluid layers: $\tau \sim \frac{\partial v_x}{\partial y}$.

At the end of the 18th century and the beginning of the 19th century, numerous and extensive investigations on the flow of pipes and open channels started to be carried out by scientists and engineers such as Du Buat, Girard, Prony, Coulomb, and especially Poiseuille (1797-1869). Being interested in the circulation of blood in capillary vessels, he then used glass capillaries of very much smaller bore that any of his predecessors. Long series of measurements of the quantity Q of liquid discharged in unit time were carried out in function of the different factors. He then obtained in the forties the relation $Q = K\frac{PD^4}{L}$ where P is the pressure, D and L respectively the diameter and the length of the capillary (Poiseuille, 1840a; Poiseuille, 1840b; Poiseuille, 1846). And in 1856, Wiedemann (1826-1899) as well as Hagenbach in 1860, deduced mathematically, by making use of Newton's hypothesis, that the constant K is such that (Wiedemann, 1856; Hagenbach, 1860)

$$Q = \frac{\pi P R^4}{8\eta L} \tag{3}$$

where R is the radius of the capillary. A new factor η appears which Wiedemann proposed to call *Zähigkeitconstante der Flüssigkeiten*, that is, the *viscosity coefficient of the liquids* (Wiedemann, 1856). So for the first time the term viscosity was used in a technical sense.

In parallel, the first scientist having theoretically used the property of viscosity in the fundamental equations of hydrodynamics was the French civil engineer Navier (1785-1836) in a mémoire read in 1822 to the "Académie des Sciences" (Navier, 1823). His idea was that any pressure tends to reduce the distances between the molecules of the fluid. Taking into account the intermolecular repulsive forces produced by this action, which gave him the explanation of the Newton's *lack of slipperiness*, he added to the pressure acting on a small volume the contribution of this effect ($P\delta_{ij} \to P\delta_{ij} - \eta\frac{\partial v_i}{\partial r_j}$), that is a dissipative term implying the presence of the viscosity coefficient η. Being rediscovered by Stokes (1819-1903) in 1845 without introducing any molecular behavior (Stokes, 1845), the history has given the name of these two persons to the famous *Navier-Stokes equations*

---

[11] In his Principia, Newton made precisely the following hypothesis:

> The lack of slipperiness of the parts of a fluid is, other things being equal, proportional to the velocity with which the parts of the fluid are separated from one another (Book II, Section 9 in Newton (1999)).

$$\rho\frac{dv_i}{dt} = -\frac{\partial P}{\partial r_i} + \eta\left(\frac{\partial^2 v_i}{\partial r_j\,\partial r_j}\right) + f_i\ . \tag{4}$$

On the other hand, in 1829, Poisson (1781-1836) derived an equation in a mémoire devoted to the motion of solid bodies and fluids in which appeared a second constant (Poisson, 1831), in addition to the viscosity introduced by Navier. These two constants measure the response of the substance to two different types of forces: the first tending to shear it, and the second tending to expand or contract it. We saw above that the first type is characterized by the viscosity coefficient η which is more precisely called *shear viscosity.* But until sixty years ago, most of the standard works did not consider the other constant η' called the *second viscosity coefficient* by making it simply proportional to the shear viscosity. Since Stokes (1845) , it was usually admitted that

$$\eta' + \frac{2}{3}\eta = 0\ , \tag{5}$$

which is generally called *Stokes' relation.* Stokes used this relation by arguing that in most cases of practical interest the rate of expansion would be so small that the value assigned to this sum would be unimportant. And before sixty years ago no direct determination of η' has yet been made. Indeed it is only in 1942 that Tisza made this determination by evaluating the ratio $\frac{\eta'}{\eta}$ when ultrasonic vibrations are passed through a liquid (Tisza, 1942). He then showed that this vanishing relation is not justified in general and that the re-introducion of the second viscosity coefficient is necessary in order to get an agreement between the theory and experiments (Markham et al., 1951; Karim and Rosenhead, 1952; Brush, 1962b). Consequently, a new coefficient ζ can be defined as

$$\zeta = \eta' + \frac{2}{3}\eta \tag{6}$$

and is the so-called *bulk viscosity coefficient*.

The bulk viscosity is more difficult to measure experimentally than the shear viscosity. Moreover, its measure is generally less known. The different ways to measure it imply the phenomenon of absorption of sound in liquids and gases. Before Tisza (1942) , it was well-known that the absorption of sound in polyatomic gases and in liquids obtained experimentally exhibited quite great disagreements with the absorption predicted by the classical theory based on viscosity. Previously, it was usually admitted that the absorption is given as

$$\alpha_{\mathrm{cl}} = \frac{2\pi^2\nu^2}{\rho v_s^3}\left[\left(\frac{1}{c_V} - \frac{1}{c_P}\right)\kappa + \frac{4}{3}\eta\right] \ , \tag{7}$$

where ν is the acoustic frequency, ρ the mass density, v_s the acoustic velocity, c_V and c_P the specific heats respectively at constant volume and constant pressure, and κ the thermal conductivity. Hence, only the shear viscosity was considered. In his paper (Tisza, 1942), he pointed out that Stokes' assumption (5) is not justified except in dilute monoatomic gases. However, for polyatomic gases and liquids, the bulk viscosity should be quantitatively quite important, sometimes more than the shear viscosity (see Table 1). Consequently, Tisza obtained a modified expression for the absorption coefficient in which the bulk viscosity is added in the expression (7) as follows

$$\alpha_{\mathrm{obs}} = \frac{2\pi^2\nu^2}{\rho v_s^3}\left[\left(\frac{1}{c_V} - \frac{1}{c_P}\right)\kappa + \frac{4}{3}\eta + \zeta\right] \ . \tag{8}$$

or by considering the ratio of the viscosity coefficients $\frac{\zeta}{\eta}$

$$\frac{\zeta}{\eta} = \frac{4}{3}\frac{\alpha_{\mathrm{obs}} - \alpha_{\mathrm{cl}}}{\alpha_{\mathrm{cl}}}\left(1 + \frac{3}{4}\left(\frac{1}{c_V} - \frac{1}{c_P}\right)\frac{\kappa}{\eta}\right) \ . \tag{9}$$

On the other hand, the tangential force acting for example on the plane xy should be related to the sliding in the two perpendicular directions composing this plane, that is $\frac{\partial v_x}{\partial z} + \frac{\partial v_z}{\partial x}$ in the direction of x, and $\frac{\partial v_y}{\partial z} + \frac{\partial v_z}{\partial y}$ in the direction of y. This is why Barré de Saint-Venant (1797-1886) combined both of them in 1843 (de Saint-Venant, 1843) and obtained more precise expressions for the pressure tensor, e.g. $P_{xx} = P + 2\eta\frac{\partial v_x}{\partial x}$ and $P_{xy} = \eta\left(\frac{\partial v_x}{\partial y} + \frac{\partial v_y}{\partial x}\right)$. It allows one to write the Navier-Stokes equations in a more complete form

fluid	$\eta \times 10^5$ Pa s	ζ/η
He	1.98	0
Ar	2.27	0
H_2	0.887	32
N_2	1.66	0.8
O_2	2.07	0.4
CO_2	1.50	1.000
Air	1.85	0.6
H_2O (liquid)	85.7	3.1
Ethyl alcohol	110	4.5
Glycerine	134.000	0.4

Table 1: Shear and bulk viscosities obtained experimentally at 300 K and 1 Atm in monoatomic, diatomic and polyatomic gases, and in liquids; values given by Thompson (1972).

as[12]

$$\rho \frac{dv_i}{dt} = -\frac{\partial}{\partial r_j}\left[P\,\delta_{ij} - \eta\left(\frac{\partial v_i}{\partial r_j} + \frac{\partial v_j}{\partial r_i} - \frac{2}{3}\delta_{ij}\,\frac{\partial v_l}{\partial r_l}\right) - \zeta\,\delta_{ij}\frac{\partial v_l}{\partial r_l}\right] + f_j\,. \quad (10)$$

When the fluid is considered incompressible ($\frac{\partial v_l}{\partial r_l} = 0$), one recovers the original Navier-Stokes equation (4). Let us also mention that these equations are valid in so far as Newton's assumption is reasonable. A wide range of fluids (water, air, etc.) follows the linear relation between the shear stress τ and the velocity gradient. But liquids composed by lengthened molecules (polymers, etc.) obey to different viscosity laws, which are called non-Newtonian.

As we will see in the next section, the 19th century and the beginning of the 20th century before the first world war has seen the decisive development of the scientific atomistic conception of matter and the will to derive macroscopic phenomena like viscosity in terms of the Newtonian microscopic dynamics. These works gave rise to the powerful kinetic theory of gases.

---

[12] For further information on the history of hydrodynamics, see Rouse and Ince (1957), Tokaty (1994), Mikhailov (1994) and Stokes (1846). A more rigorous derivations of Navier-Stokes equations can be found for example in (Landau and Lifshitz, 1959).

4 Kinetic theory of gases

The idea of atomism goes back to the fifth century BC with the ancient Greek philosophers Leucippus and Democritus. However, the history of the kinetic theory of gases does not really begin until the seventeenth century when Torricelli, Pascal (1623-1662) and Boyle first established the physical nature of the "sea" of air that exerts mechanical pressure on surfaces in contact with it. This development of the concept of air pressure can be considered as part of the change in scientific attitudes which led to the explanations based simply on matter and motion instead of postulating "occult forces" or teleological principles. Consequently, Boyle (1662) discovered the law $PV = \text{const}$ now called *Boyle law.*

Usually we consider Daniel Bernoulli as the first scientist having proposed a kinetic theory of gases. Actually, in his famous treatise *Hydrodynamica* (Bernoulli, 1738), he gave a derivation of the gas law of Boyle and Mariotte from "billiard ball" model, assuming that the gas consists of a very large number of small particles in rapid motion. Moreover, using the principle of conservation of *vis viva* he concluded that pressure is proportional to the square of the velocities of gas particles, and thus proportional to the temperature. In other terms heat is nothing but atomic motion. However, in spite of these results, Bernoulli's model was forgotten. He was about a century ahead of his time with his theory. The latter success of Lavoisier's caloric theory buried it by proposing another conception of the matter based on the equilibrium between the caloric repulsion (due to the atmosphere of caloric whose density increases with temperature) and the gravitational attraction between matter particles.

In chemistry, the atomic theory emerged at the beginning of the 19th century with Dalton (1766-1844). Thanks to the laws of chemical discontinuities, that is the law of *constant proportions*[13] discovered in 1806 by Proust (1754-1822) and the law of *multiple proportions*[14] in 1802-1804 by Dalton himself, he proposed in 1808 a discrete view of matter, which should be composed by indivisible entities: the atoms which are characterized by different *atomic weights.* But these laws and experiences at that time did not reject a concept of *equivalents* first proposed in 1792 by Richter (1762-1807). Because of different ambiguities amongst the partisans of the atomic theory, the equivalentism had a great success during the 19th century. However, many new phenomena and laws were discovered that only the atomic hypothesis could explain. Among these, the law of Gay-Lussac (1778-1850) discovered in 1809[15],which allowed to Avogadro (1776-1850) to emit his famous hypothesis (1811) saying that all gases considered in the same conditions of temperature and pressure contain, for equal volumes, the same number of molecules[16]. We can also cite the law of Dulong (1785-1838) and Petit (1791-1820)[17] (1819); the emergence of structural chemistry, in particular the development of the theory of valence by Kekulé (1829-1896), organic chemistry and the concept of isomerism, and stereochemistry. But one of the most important works having played a fundamental role in the development of the atomic theory and its acceptance is the establishment of the periodic table of the elements in 1869 by Mendeleev (1834-1907) (Mendeleev, 1869a; Mendeleev, 1869b). Indeed, by assuming his periodic law in the properties such as the valence of elements when these are listed in order of atomic weight, he predicted the existence of unknown elements corresponding to the gaps in his famous table and to which a given atomic weight is associated (Brush, 1996). The discovery of some of them in the following years confirmed the periodic property of elements. In the polemic between atomists and equivalentists, Mendeleev's brilliant idea heavily contributed to the acceptance of the atomic conception of the matter amongst the chemists[18].

---

[13] The proportion between two combining elements cannot vary continuously.

[14] When two elements can combine according to different ratios, the ratios of their weight in the different cases are simple, that is integer numbers.

While the caloric theory was being brought to its final stage of perfection by Laplace, Poisson, Carnot and others, Herapath (1790-1868) in 1820 and 1821, as well as Waterston (1811-1883) in 1845 proposed their kinetic theory[19]. However both of them were rejected by the Royal Society. It must be emphasized that what needed to be established was not simply a connection between heat and molecular motion, for that was already admitted by many scientists and was not considered incompatible with the caloric theory; it was rather the notion that heat is *nothing but* molecular motion, and the idea that molecules *move freely through space* in gases rather than simply vibrating around fixed positions. This statement could not yet be accepted.

---

[15] The ratios of volumes of reacting gases are integer numbers.

[16] Let us notice that Avogadro was the first scientist having used the term *molecule* in the modern sense. The word is derived from the french word *molécule* meaning "extremely minute particle" which comes from the Latin word *molecula*. This term is the diminutive of *mole* meaning "mass, cluster, great number, multitude". On the other hand, the word *atom* is derived from the Latin term *atomus* used by Lucretius, and from the Greek word $\alpha\tau o\mu o\varsigma$, which means uncut.

[17] The atomic weight multiplied by the specific heat of an element is independent of the nature of the element.

[18] For further information on the history of atomic theory in chemistry, see for example Pullman (1995) and Leicester (1956).

[19] For further informations on their theories, see respectively Brush (1957a, 1957b).

During the period between 1842 and 1847 the general scientific and intellectual climate implied quite simultaneous works by different scientists on a concept which is one of the most important reasons of the revival of kinetic theory: the *conservation of energy.* Indeed, the influence of the German romanticism and *Naturphilosophie*, especially the idea that there must be a single unifying principle underlying all natural phenomena (Brush, 1967). Furthermore, this period inherited different discoveries of various conversion processes[20] and gave birth to this principle usually attributed to Mayer (1814-1878) (Mayer, 1842) and Joule (1818-1889) (Joule, 1847). Mayer emphasized the philosophical generality of the principle while Joule provided the experimental verification in particular cases. From that moment heat, mechanical work, electricity and other apparently different entities are considered as different forms of the same thing, now called *energy.* And in 1847 Joule and Helmholtz (1821-1894) indicated quite clearly that *mechanical* energy is regarded as the basic entity. It was this prejudice toward mechanical explanations that made the kinetic theory appear to be an obvious consequence of the principle of conservation of energy. If we are convinced that heat and mechanical energy are interconvertible, what is more natural than to conclude that heat is a mechanical energy?

The real breakthrough for the kinetic theory took place when Krönig (1822-1879) assumed in 1856 (Krönig, 1856) that the molecules of gas move with constant velocity in straight line until they strike against other molecules, or against the surface of the container[21]. But the kinetic theory was still confronted to objections like those supported by Buys-Ballot (1817-1890). Since the kinetic theory claimed that the velocities of molecules were of the order of several hundred meters per second, he pointed out that we would expect two gases to mix with each other very rapidly. However, the experience shows that the process takes a certain time, of the order of several minutes. In order to answer this objection he showed that, in real gases, the molecules could not travel without colliding with other molecules. Consequently, Clausius (1858) introduced a new concept: the so-called *mean free path* of a molecule between two successive collisions. This concept was major not only for the further developments, but also because it establishes in concrete terms one of the most fundamental ideas of the kinetic theory of gases rejected in the past: that molecules can move freely through space and yet collide with each other.

---

[20] Oersted's discovery of electromagnetism (1820), Seebeck's discovery of thermoelectricity (1822), Faraday's many discoveries in electricity and magnetism, and many others.

[21] Actually his publication did not represent a real advance compared to previous works by Bernoulli and Herapath. But his influence in the physics community was important and induced an special interest among the physicists for the kinetic theory.

The early kinetic theorists assumed that molecules tending to equilibrium move all at the same velocity. Maxwell (1831-1879) was the first scientist who introduced the idea of random motion for the molecules – hence statistical considerations[22]. In 1860, in his first paper on kinetic theory entitled *Illustrations of the dynamical theory of gases* (Maxwell, 1860), he suggested that, instead of tending to equalize the velocities of all molecules, the successive collisions would produce a statistical distribution of velocities in which all might occur, with a known probability. For thermal equilibrium he could then derive from symmetry considerations his famous distribution function which in modern notation is given $f(v) = 4\pi(m/2\pi k_B T)^{3/2}\ v^2\ \exp(-mv^2/2k_B T)$. Hence statistical character appeared to be a fundamental element of kinetic theory.

In 1859, Maxwell came to the kinetic theory as an exercise in mechanics involving the motions of systems of particles acting on each other only by impact. Being interested in viscosity he proposed a mechanism which allowed him to establish a relation between the (shear) viscosity and the mean free path. By considering gas divided into parallel layers and supposing that the motion is uniform in each layer but varying from one to another, he showed that the viscosity η should have to be proportional to the mean free path $\langle l \rangle$, the mass density ρ and the molecular mean velocity $\langle v \rangle$, relation written as

$$\eta_M = \frac{1}{3}\ \langle v \rangle\ \rho\ \langle l \rangle\ . \tag{11}$$

He also published this result in his first article on the kinetic theory (Maxwell, 1860). However, as Clausius and himself concluded, the mean free path is inversely proportional to the density, which implied the surprising result that the viscosity in a gas does not depend on its density. Whereas it was known that this dependence does exist in liquid, and according to the opinion of Stokes, Maxwell thought his predictions were absurd and therefore that the kinetic theory is wrong, or at least inadequate. But at that time a few experiments on the gas viscosity had already been realized. The first studies probably began in the 1840's with e.g. Graham (1846, 1849). However, accurate experiments had not yet been done on the viscosity of *gases*, therefore, in 1866,

---

[22] Actually Maxwell was influenced by works on statistics, in particular the works realized by Quetelet (1796-1874) on the height distribution of a population of soldiers. Quetelet stated the idea of the 'average man' with 'average height' etc., as the result of an experimentally established normal distribution function. Maxwell worked out the idea of an 'average molecule' in analogy with that of the 'average man'.

Maxwell himself carried out his own experiment and found out that viscosity remains constant over a large range of pressure (Maxwell, 1866). This work played an important role in the development of the kinetic theory and its acceptance by most of scientists who so far had been doubting it. For instance, in 1865, Loschmidt (1821-1895) gave the first convincing estimate of the diameter of an air molecule (about 10 Å which is about four times too large) as well as the Avogadro number (about 6.025×10^{23}) (Loschmidt, 1865). On the other hand, Maxwell formula, combined with the equation of state for real gases derived in 1873 by van der Waals (1837-1923)[23] (van der Waals, 1873), belongs to the list of thirteen different phenomena mentioned in 1913 by Perrin (1870-1942) and allowing one to evaluate the atomic magnitudes as well as the Avogadro number (Perrin, 1991).

As the kinetic theory establishing a microscopic mechanical basis to the macroscopic processes such as viscosity was attracting more and more attention by physicists, the idea to reconcile the second law of thermodynamics with the principles of mechanics emerged and the first scientist having been concerned by it was Boltzmann (1844-1906). His first major achievement (Boltzmann, 1868) was to extend Maxwell's distribution law to the case of an external force field (field deriving from the potential energy V) is present. He obtained the so-called *Boltzmann factor* which, combined with Maxwell's velocity-distribution law, constitutes the basic principle of statistical mechanics (Maxwell-Boltzmann distribution law).

In 1872, Boltzmann attempted to establish an equation describing the changes in the distribution resulting from collisions between molecules. Indeed he considered the single particle distribution function $f(\mathbf{r}, \mathbf{v}, t)$ so that $f(\mathbf{r}, \mathbf{v}, t)\delta\mathbf{r}\delta\mathbf{v}$ gives the average number of molecules in the infinitesimal volume $\delta\mathbf{r}\delta\mathbf{v}$ around the position $\mathbf{r}$ and the velocity $\mathbf{v}$. The evolution of this distribution function is governed by the integro-differential equation which is now called the *Boltzmann equation* (Boltzmann, 1872; Boltzmann, 1995)

$$\frac{\partial f}{\partial t} = -\mathbf{v} \cdot \frac{\partial f}{\partial \mathbf{r}} + J_B(f, f) \; . \tag{12}$$

Here J_B is the binary collision term[24] taking only two particle collisions into account, which is a good approximation for dilute gases. An important assumption made by Boltzmann is the so-called *Stosszahlansatz*[25], which assumes that the velocities of colliding particles must be uncorrelated. Later Jeans (1877-1946) developed this statement and introduced the assumption of *molecular chaos*[26] (Jeans, 1903; Jeans, 1904). Boltzmann showed that collisions always push f toward the equilibrium Maxwell distribution. In particular the quantity $H = \int f \log f$ always decreases with time unless f is the Maxwell distribution in which case H remains constant. This is *Boltzmann's H-theorem* $\frac{dH}{dt} \leq 0$ (Boltzmann, 1872). Boltzmann suggested that H could be considered as a generalized entropy having a value for any state, contrary to the thermodynamic entropy defined only for equilibrium states. On the other hand, he explained the reason why the Maxwell distribution law is the one corresponding to the thermal equilibrium by showing that this distribution is the one most likely to be found, because it corresponds to the largest number of microstates. By comparing the two approaches (the kinetic approach and the one based on probabilities), Boltzmann could conclude that the process of irreversible approach to equilibrium, which is a typical example of entropy-increasing process, corresponds to a transition from less probable to more probable microstates. Entropy itself can therefore be interpreted as a measure of probability. By defining Ω the probability of a macrostate, which is proportional to the number of associated microstates, Boltzmann proposed a generalized entropy of a macrostate $S = k_B \log \Omega$, now called the *Boltzmann entropy.*

---

[23] Work based on the *virial theorem* introduced by Clausius (1870): "the mean kinetic energy of the system is equal to the mean of its virial multiplied by $-\frac{1}{2}$: $\overline{E} = -\frac{1}{2}\sum_i \overline{\mathbf{r}_i \cdot \mathbf{F}_i}$", where $\mathbf{r}_i$ and $\mathbf{F}_i$ are respectively the position of the ith particle and the force acting on it.

[24] $J_B(f, f) = \int \sigma_{\text{diff}} |\mathbf{v}_1 - \mathbf{v}| \, (f' f'_1 - f f_1) \; d\mathbf{v}_1 d\Omega$ where f and f_1, f' and f'_1 are the functions of the two particles respectively before and after the collision. σ_{diff} represents differential cross section and ω the angular variables of integration. For a detailed derivation of the Boltzmann equation, see . Boltzmann (1995) and Dorfman (1999).

With the turning of the century Gibbs (1839-1903) transformed the kinetic theory into a more abstract mathematical theory based on the concept of ensembles of similar systems. Actually Boltzmann (1884) himself was working in this way by introducing the so-called *Ergoden*, which corresponds to the *microcanonical ensemble* of Gibbs (1839-1903). In his definitive treatise, Gibbs (1902) presented this ensemble as one of a series of ensembles. This new formalism appeared to be a powerful technique for calculating and predicting the physical properties of bulk matter *at equilibrium* from the properties of atoms.

Despite all the brillant discoveries due to the atomic hypothesis of matter accumulated at the end of the 19th century, a lot of scientists (amongst the most important) continued to reject the atomic theory. In general, they belonged to movements called *equivalentism* represented by especially Berthelot (1827-1907), *energetics* with Ostwald (1853-1932) and Duhem (1861-1916), or *empiriocritism*, the doctrine of Avenarius (1843-1896) and Mach (1838-1916)[27]. The debate then were still vigorous between anti-atomists and the partisans of the kinetic theory, especially Boltzmann. Fortunately, during the first decade of the 20th century, the triumph of the kinetic theory and the atomic hypothesis arrived with the work done in 1905 by Einstein (1879-1955) on the *Brownian motion* (Einstein, 1905). Being aware of the attacks on kinetic theory by Mach, Ostwald and others, Einstein started his article by contrasting the predictions made by thermodynamics and by the kinetic theory (in particular by pointing out that thermodynamics distinguishes the Brownian particle and the "hypothetic" molecules composing the liquid, whereas kinetic theory does not and says that the colloidal particle should follow the Clausius' *equipartition theorem*). By combining Stokes' formula for the force on a sphere moving through a viscous fluid, and the formula for the osmotic pressure of dissolved molecules, he derived an expression for the mean-squared displacement of the Brownian particle submitted to the molecular agitation, which gave a way to evaluate the Avogadro's number N_{Av}. In 1908, Perrin (1870 - 1942) made different experiments in application to the Einstein's Brownian-motion theory and found $N_{Av} = 6.7 \times 10^{23}$ (Perrin, 1908). Later, in 1913, he published his now famous book *Les atomes* (Perrin, 1991) in which he proposed thirteen different experiments based on different phenomena for evaluating Avogadro's number. The quite narrow range $6.0 - 7.5 \times 10^{23}$ thus obtained is such that we now attribute the demonstration of the discrete feature of the matter to Perrin, especially due to his efforts to convince the opponents of atoms' existence[28]. Whereas Ostwald and others recognized it, some of them like Mach and Duhem continued to reject them (Pullman, 1995).

---

[25] "Assumption about the number of collisions".

[26] For some informations on the distinctions between both assumptions, see (Ehrenfest and Ehrenfest, 1990).

If the performances made by the kinetic theory were already numerous, new works and predictions were added and thus confirmed the depth of this theory. In 1911, Enskog (1884-1947) predicted that diffusion would appear in a mixture when one imposes a temperature gradient, phenomenon now called *thermal diffusion*. One year later, Chapman (1888-1970) derived an expression for the thermal diffusion coefficient.

Around the end of the 19th century and the beginning of the 20th century different works based on the mean-free-path method were achieved by a lot of physicists (Maxwell, Boltzmann, Tait, Rayleigh,...)[29] to improve the expression for the viscosity in low-density gases. But one has to wait for the almost simultaneous works carried out by Chapman in 1916 and Enskog in 1917 to obtain an expression for the viscosity derived from the kinetic equation[30] (Chapman, 1917; Enskog, 1917). The derivation known today is that of Enskog, and is based on a series solution of Eq.(12) for the distribution function by introducing a parameter λ. He therefore derived analytical expressions for the transport coefficients for general potential of interactions. In the case of the hard-sphere potential, the viscosity coefficient is written as

---

[27] For further informations, see for example Pullman (1995), Brush (1967) and Kubbinga (1990).

[28] Let us mention that the recent measurements gives us $6.02214199(47) \times 10^{23}$ for the current value of the Avogadro's number (Mohr and Taylor, 1998).

$$\eta_B = 1.0162\,\frac{5}{16\,\sigma^2}\sqrt{\frac{mk_BT}{\pi}} \tag{13}$$

where σ and m are respectively the diameter and the mass of the particles. This expression for the viscosity is now called the *Boltzmann viscosity.* This new expression confirms the temperature-dependence and the absence of density-dependence predicted by earlier theory, especially that of Maxwell in Eq.(11)[31]. Furthermore, being available for any potential interaction, the Chapman-Enskog's developement turned out to be a method for determining the actual force law of real molecules by comparing the experimental and theoretical results for transport properties as viscosity and thermal conductivity. This was the way followed by Lennard-Jones (1894-1954) until the early thirties when he proposed the now called *Lennard-Jones 6-12 potential* which is still considered today as the most realistic potential existing between particles (Lennard-Jones, 1931)[32].

Maxwell and others showed both theoretically and experimentally that the viscosity coefficient is independent of the density in dilute gases. However, even in moderately dense gases, it appears that this property is no longer observed. In 1922, Enskog proposed an extension of his previous method in order to predict this dependence (Enskog, 1922). Considering a hard-sphere system he assumed that the collision rate in a dense gas would be changed by a factor χ which could be related to the equation of state. He modified the Boltzmann equation (12) by replacing ff_1 in which both f and

---

[29] Especially the dependence in temperature and the factor $\frac{1}{3}$ in Eq.(11).

[30] Notice that Chapman started from the Maxwell's transport theory whereas the Enskog's derivation was based on Boltzmann's. For a brief discussion on both methods, see Brush (1962a).

[31] By substituting the mean velocity $\overline{v}$ and the mean free path $\overline{l}$ by their expression in terms of temperature and atomic properties, one obtains $\eta = \frac{2}{3\,\sigma^2}\sqrt{\frac{mk_BT}{\pi^3}}$.

[32] For more historical elements of the development of expression for interatomic forces, see for example Brush (1970).

f_1 are evaluated in the same point in space, by $f(x, y, z, ...)f_1(x_1, y_1, z_1, ...)$, where the point points (x, y, z) and (x_1, y_1, z_1) must be separated by the distance σ. In such a way, he obtained[33]

$$\eta_E = \eta_B \left(\frac{1}{\chi} + \frac{4}{5}\, b_0 n + 0.7614\ b_0^2 n^2 \chi \right) \tag{14}$$

in which the factor $\chi = 1 + \frac{5}{8} b_0 n + 0.2869\ (b_0 n)^2$, $b_0 = \frac{2\pi\sigma^3}{3}$ and n the number density. A density-dependent expression for the viscosity is therefore obtained[34]. One had to wait for a decade to compare Enskog's theoretical predictions to experimental data. Indeed, Michels and Gibson (1931) measured the viscosity of nitrogen gas at pressure up to 1000 atm (1013250 hPa) and obtained a good agreement with the semi-empirical Enskog theory presented below. Using this method, they also evaluated the dimensions of the gas molecule at different temperatures. Let us point out that the only work allowing Enskog to test his theory was the one accomplished by Warburg and von Babo (1882) for carbon dioxide. The agreement is quite good (see Table 2).

In the fifties and sixties, new experimental data for noble gases (for viscosity as well as for thermal conductivity) were added to allow a comparison and a test of theoretical predictions[35]. The particular advantage of Enskog's theory compared to other theories for dense gases is that it requires the adjustment of only a few parameters like the diameter of the spheres in order to observe an agreement between Enskog's viscosity and the experimental data for a larger range of density[36].

---

[33] For the development of Enskog's theory, see for example Chapman and Cowling (1970) and Hirschfelder et al. (1954).

[34] Let us mention that, in 1899-1900, by considering the analogy with the modification of the equation of state for ideal gas when the effect of finite molecular size is taken into account, and the correction the mean free path due to the effect of excluded volumes on the collision rate, Jäger (1865-1938) modified the dilute gas viscosity formula (11) and obtained

$$\eta_J = \eta_M \left(\frac{1}{A} + 8\ b_0 n + 16\ b_0^2 n^2 A \right)$$

where $A = 1 + \frac{5 b_0 n}{2} + \cdots$. The similarity with the Enskog viscosity (14) is remarkable (Brush, 1976).

[35] see for example Senger (1965, 1968), Hanley et al. (1972).

[36] Indeed effective values may be attributed to the parameters b and χ. It was Michels and Gibson (1931) who introduced such a procedure by identifying the pressure P in the equation of state for rigid spheres with the thermal pressure $T(\partial p/\partial T)_V$ of the real gas

$$bn\chi = \frac{1}{R}\left(\frac{\partial PV}{\partial T}\right)_V - 1\ .$$

Mass density	Kinematic viscosity (experimental)	Kinematic viscosity (calculated)
0.100	1.80	1.91
0.170	1.12	1.17
0.240	0.908	0.899
0.310	0.784	0.774
0.380	0.724	0.719
0.450	0.702	0.701
0.520	0.704	0.705
0.590	0.722	0.724
0.660	0.756	0.754
0.730	0.795	0.792

Table 2: Kinematic viscosity $\frac{\eta}{\rho}$ $[10^3 poises/(g/cm^3)]$ versus the mass density $[g/cm^3]$ at temperature $T = 313.45K$, near the critical point. Comparison between the experimental results obtained by Warburg and von Babo (1882) for carbon dioxide and Enskog's predictions. As the density increases, he kinematic viscosity goes through a minimum after which it increases. Enskog's predictions confirm this property (data given in Enskog (1922)'s paper).

In Table 1, it is shown that the bulk viscosity is vanishing in the dilute monoatomic gases (argon and helium) as predicted by the Enskog theory. Indeed Enskog's bulk viscosity (15) for hard spheres

$$\zeta = 1.002\chi\eta_B(b\,n)^2 \tag{15}$$

vanishes for very low density. However, a non-vanishing ratio ζ/η has been measured in liquid argon (Naugle and Squire, 1965) as well as in dense gaseous argon (Madigosky, 1967). A comparison with the modified Enskog theory briefly presented above can be found in Hanley and Cohen (1976). Consequently Stokes' relation is not shown to be justified against many cases, not only through theoretical predictions, but also experimentally.

For low densities, one should require that the the Enskog viscosity (14) reduces to the Chapman-Enskog viscosity (13) by requiring $\lim_{n\to 0}\chi = 1$, so that we get (Hirschfelder et al., 1954)

$$b = B + T\frac{dB}{dT}$$

where B is the second virial coefficient.

In the sixties it was observed that transport coefficients, in particular the viscosity coefficient η, cannot be expressed in a power series in terms of the density n ($\eta = \eta_B + \eta_1 n + \eta_2 n^2 + \eta_3 n^3 + \dots$) Indeed it has been shown that correlations between molecules is observed over large distances, larger than the range of the intermolecular interaction. Hence Dorfman and Cohen (1967) obtained theoretically that the coefficient of the quadratic term contains a contribution proportional to the logarithm of the density so that the previous density expansion should be rewritten as

$$\eta = \eta_B + \eta_1 \, n + \eta_2^{'} \, n^2 \ln n + \eta_3 \, n^3 + \dots \; . \tag{16}$$

Nevertheless although the importance of the discovery of long-distance correlations is great in the modern kinetic theories, their contributions do not seem to be important, as it was showed by Sengers (1966) for the viscosity in a hard-disk gas , and later by Kamgar-Parsi and Sengers (1983) in a gas of hard spheres . On the other hand, since the early seventies several attempts have been made to detect this logarithmic-density dependence of experimental data, in particular for the viscosity coefficient[37]. Even if the viscosity is generally the transport coefficient measured with the highest precision, it has never been shown that the addition of the logarithmic term was necessary. Although we now know that Enskog viscosity is not exactly correct, the latter may still be used in order to compare numerical results. This is why, despite these facts, we consider Enskog's theory in this thesis[38].

Following Enskog theory proposed in 1922, we have to wait 1946 to observe a revival of interest for the extension of kinetic theory to dense gases and liquids. Indeed, after an original idea of Yvon (1935), a hierarchy of coupled equations ruling the time evolution of the reduced n-particle distribution functions was derived by Bogoliubov, Born, Green, Kirkwood and Yvon such that, by truncating this BBGKY hierarchy, kinetic equations were derived leading to expansions of the transport properties in terms of the particle density (Yvon, 1935; Bogoliubov, 1946; Born and Green, 1946; Kirkwood, 1946).

In 1928, Pauli (1900-1958) derived a *master equation* in order to apply quantum mechanics to irreversible processes (Pauli, 1928). This derivation also contained an assumption analogue to Boltzmann's *Stosszahlansatz* – the repeated random phase assumption. Since then, different works have been completed by means of master equations, such as the generalization of the classical Boltzmann equation to quantum systems by Uehling and Uhlenbeck (1933). In the fifties, master equations were derived for weakly coupled systems by van Hove (1955, 1957, 1959) as well as Brout and Prigogine (1956). Kinetic equations were also obtained for plasmas since thanks to Landau (1936) and Vlassov (1938). In this context, Balescu (1960) and Lenard (1960) derived in the sixties the so-called *Lenard-Balescu* transport equation for plasmas.

---

[37] See for example Kestin et al. (1971, 1980, 1981).

[38] A historical survey of the discovery of the logarithmic-density dependence is given by Brush (1972), pp.72-80.

In parallel, the relationship between molecular fluctuations in systems due to thermal agitation, and transport coefficients characterizing the irreversible processes was studied starting with Einstein's work on the Brownian motion (Einstein, 1905), giving the self-diffusion of the Brownian particle in terms of the variance of its position

$$D = \lim_{t\to\infty} \frac{\left\langle [x(t)-x(0)]^2 \right\rangle}{2\,t} \;, \tag{17}$$

to reach *the linear response theory* developed in the fifties by Green (1951, 1960), Kubo (1957), and Mori (1958a). This theory expresses the transport coefficients in terms of the integral of autocorrelation function. In particular, the viscosity coefficient is written as the autocorrelation of the microscopic current $J(t) = \sum_a \frac{1}{m} p_{ax} p_{ay} + \frac{1}{2} \sum_a \sum_{b\neq a} F_x(\mathbf{r}_{ab})(r_{aby})$

$$\eta = \lim_{V\to\infty} \frac{1}{V k_B T} \int_0^\infty \langle J(0) J(t) \rangle \; dt \;. \tag{18}$$

Moreover, aiming at establishing an Einstein-like relation for transport processes, Helfand (1960) derived formulas relating the transport coefficients to the variance of their associated quantities now called *Helfand moment* $G(t)$. In particular, in the case of viscosity, such relationship is written as

$$\eta = \lim_{V\to\infty} \lim_{t\to\infty} \frac{\left\langle [G(t)-G(0)]^2 \right\rangle}{2tVk_BT} \quad ; \quad G(t) = \sum_{a=1}^{N} p_{ax}(t) y_a(t) \;, \tag{19}$$

and has the same structure than Eq. (17). On the other hand, it can be shown that the last two relations are equivalent by noting that $J(t) = \frac{dG(t)}{dt}$.

The transport coefficients are usually calculated based on some kinetic equation. Each kinetic equation involves a stochastic assumption such as Boltzmann's *molecular chaos* which is reached by some truncation of the evolution equation. Such phenomena as the transport processes, playing such an important role in nature, must present a more fundamental justification in terms of the intrinsic properties of the underlying microscopic dynamics. We can conclude that microscopic chaos seems to be the key for a better understanding of the irreversibility of the macroscopic phenomena.

5 Theory of chaos

It is interesting to note that the idea of considering a gas as a disordered state is old. During the 16th century Paracelsus (1493-1541) studied gases and observed that the air and vapors did not have a fixed volume or shape. He then gave them the name of *chaos* in analogy with $\chi\alpha o\varsigma$ in the greek mythology, being at the origin of the universe. Later, van Helmont (1577-1644), who is the first to have made the distinction between the different gases and the air, invented the word *gas* according to the flemish pronounciation of chaos (Leicester, 1956).

In the end of the 19th century was generally accepted that a large number of degrees of freedom is a necessary condition to observe unpredictable behavior. This great number was the necessary element in order to introduce a statistical description of phenomena having a deterministic underlying dynamics. However, in 1892, Poincaré (1854-1912) showed that a low-dimensional deterministic system, namely the *three-body problem*, exhibits an unpredictable behavior which is now commonly called *chaos* (Poincaré, 1892). Since his work, a large number of nonlinear dynamical systems, the time evolution of their variables $\mathbf{x}$ being given by $\frac{dx_i}{dt} = F_i(x_j)$, have been discovered. In these systems, chaos appears to be a generic phenomenon rather than an exotic one (Nicolis, 1995).

As a matter of fact, in 1873, Maxwell himself had already emphasized the sense of unstable behaviors

> When the state of things is such that an infinitely small variation of the present state will alter only by an infinitely small quantity the state at some future time, the condition of the system, whether at rest or in motion, is said to be stable; but when an infinitely small variation in the present state may bring about a finite difference in the state of the system in a finite time, the condition of the system is said to be unstable. It is manifest that the existence of unstable conditions renders impossible the prediction of future events, if our knowledge of the present state is only approximate, and not accurate [39].

On the other hand, in the late years of the same century, Hadamard (1865-1863) studied the geodesic flows on a negative-curve surface and concluded that

---

[39] Cited by Hunt and Yorke (1993). The text of the conference of Maxwell may be found in Garnett (1882).

> each change of the initial direction, as small as possible, of a geodesic which remains at a finite distance is enough for implying an arbitrary variation of the behavior of the trajectory (Hadamard, 1898).

By considering Hadamard's work in a book, Duhem (1906) also concluded in the chapter entitled *Exemple de déduction mathématique à tout jamais inutilisable* that, in Hadamard's billiard, the trajectory obtained mathematically becomes unusable for a physicist since the an experimental measure realized by any physical procedure as precise as possible is always endowed of an error which grows with time. Furthermore, Poincaré clearly explained that

> A very small cause which escapes our notice determines a considerable effect that we cannot fail to see, and then we say that the effect is due to chance (Poincaré, 1908),

implying that determinism and chance may be combined thanks to the unpredictability. The only mathematician having taken into account what we now call the *sensitivity to the initial conditions*, property put in evidence by Poincaré and the others, was Birkhoff (1884-1944) who developed in the twenties the theory of *billiards* (Birkhoff, 1927). Otherwise, one had to wait for the works done in particular by Lorenz (1963) in meteorology to consider again the *sensitivity to the initial conditions* defining chaos, and the *attractor* he put in evidence. Others like Smale (1967) considered global behavior of phase space rather than a particular trajectory, and invented his *horseshoe* (see Fig.1) in order to get a visual analogy to the sensitive dependence to the initial conditions. Finally, Feigenbaum (1978) discovered the universality of the *period-doubling bifurcation cascade*[40].

In 1892 Lyapunov (1857-1918) defended his doctoral thesis *The general problem of the stability of motion* (Lyapunov, 1892). In his work he proposed a method which provides ways of determining the stability of sets of ordinary differential equations. He introduced the well-known *Lyapunov exponents*, quantities characterizing the growth ($\lambda_i > 0$) or decay ($\lambda_i < 0$) rate of the distance between two nearby trajectories in the phase space of dynamical systems (Eckmann and Ruelle, 1985). The positivity of such quantities expresses a dynamical instability and induces a *sensitivity to initial conditions*, property defining *chaos*. Two trajectories initially very close separate exponentially in time. This sensitivity to initial conditions limits the possible predictions on the trajectories because they are only known through a given precision $\epsilon_{\text{initial}}$. By considering a rate of separation of very close trajectories given by the maximum Lyapunov exponent $\lambda_{\max}$, the error between the predicted and the actual trajectories grows as $\epsilon_t \simeq \epsilon_{\text{initial}} \exp(\lambda_{\max} t)$. After a finite time, the error becomes larger than the final allowed precision ϵ_{final}, which then defines the Lyapunov time $t_{\text{Lyap}} \simeq (1/\lambda_{\max}) \ln(\epsilon_{\text{final}}/\epsilon_{\text{initial}})$. Given these initial and final precisions, predictions after the Lyapunov time are no longer relevant. This result is the requirement of going a statistical description.

---

[40] For further informations see Gleick (1987) and Ott (1993).

Inspired by the ideas of Krylov (1979), who argued that trajectories in phase space for simple fluid separate exponentially, Sinai and his coworkers proved the ergodic hypothesis for billiard systems, and characterized the stochastic-like behavior of these deterministic systems (Sinai, 1970). This type of stochastic-like behavior was then studied extensively for simple systems with few degrees of freedom.

The unstable character of dynamical systems is therefore such that even deterministic systems can generate random behaviors. But on the other hand, this instability of the dynamics produces information in time allowing to reconstruct the system trajectory in phase space. Indeed the separation in time of nearby trajectories gives us the possibility to distinguish the trajectories. In this context, in the late fifties Kolmogorov and Sinai applied the concept of entropy per unit time introduced a decade before in the information theory by Shannon to the dynamical systems. He defined the so-called *Kolmogorov-Sinai entropy* h_{KS} (Eckmann and Ruelle, 1985). This



Figure 1: *Smale's horseshoe*: Succession of stretchings and foldings illustrating the evolution of phase-space due to the chaotic dynamics.

new quantity measures the (exponential) rate at which information is obtained in time in random processes[27]. The KS entropy per unit time is the dynamical analogue of the entropy per unit volume defined in equilibrium statistical mechanics. Later, in the seventies, Sinai, Bowen, and Ruelle proposed the mathematical foundations of a new formalism for chaotic systems applying the techniques used in statistical thermodynamics, which is called the *thermodynamic formalism* (Ruelle, 1978; Beck and Schlögl, 1993). In this formalism, a new quantity, the *topological pressure*, was defined and played a role in dynamical systems very similar to that of the free energy for statistical-mechanical systems.

The two aspects of chaos, that is the *dynamical instability* characterized by the Lyapunov exponents, and the *dynamical randomness* by the KS entropy are strongly related, the second one being the consequence of the first one. In 1977, Pesin proposed a theorem for closed systems (systems without any escape of trajectories out of the initial phase space) which is now known under the name of *Pesin's identity*, which relates the KS entropy to the sum of all the positive Lyapunov exponents of the system (Pesin, 1977; Eckmann and Ruelle, 1985)

$$h_{\mathrm{KS}} = \sum_{\lambda_i > 0} \lambda_i \ . \tag{20}$$

As we saw, the instability of the dynamics in phase space induces an exponential separation of trajectories characterized by the Lyapunov exponents. Because the phase-space volume accessible to the trajectories is finite (e.g. the volume defined by the energy of the system), they have to fold onto themselves. We then observe successive stretchings and foldings of phase-space volumes, as the Smale's horseshoe illustrates it (see Fig.1), creating strange objects (after an infinite number of such operations) nowadays called *fractals*.

Whereas the term *fractal* is associated with Mandelbrot, the history of this intriguing discipline of mathematics began during the 1870's, when continuous functions without derivatives were discovered. For a long time, the idealization of nature implied a smooth and regular representation of real objects (Chabert, 1994). In mathematics, one dealt with continuous functions such that a tangent could be drawn at (almost) each point. However, Riemann (1826-1866) already claimed a contradictory opinion. And in the 1870's Weierstrass (1815-1897) gave an example having no derivative in any point[41]. In geometry, mathematicians, such as Koch (1870-1924) in 1904,

---

[41] The paper was read in 1872 in the Royal Prussian Academy of Sciences, but was only



Figure 2: *Koch's curve*: the first four steps of construction. The operation consists in dividing each segment into 3 parts, and on the one in the medium is constructed an equilateral triangle.

proposed continuous curves without a tangent at any point obtained by an elementary geometric construction (Koch, 1904). *Koch's curve* depicted in Fig.2 is a clear example of such objects. Moreover, the Japanese mathematician O. Takagi (1875-1960) working at Göttingen proposed in 1903 a simple example of continuous but nondifferentiable function known today as *Takagi's function* (Takagi, 1903). On the other hand, studying the Brownian motion, Perrin observed experimentally that the trajectory drawn by the Brownian particle is highly irregular and he deduced that

> If the functions with derivative are the simplest ones, the simplest to be treated, functions with derivatives are the exception; or, to use geometric language, curves with no tangent at any point become the rule (from Ref. (Perrin, 1991)).

Indeed according to the precision of the measure, the length of the trajectory is different, the larger the precision is, the larger is the trajectory. The relevance of the notion of length thus vanishes and has to be replaced by a new quantity characterizing such objects, that is the *dimension.* In this context, Hausdorff (1868-1942) proposed in 1919 a new notion of dimension which is no longer a whole number, but can take noninteger values (Hausdorff, 1919). For example, in the case of Koch's curve, its *Hausdorff dimension* is neither equal to one, that is the dimension of a line, nor equal to two, the dimension of the plane, but between both of them, that is 1.26 .

By introducing the term *fractal*[42], Mandelbrot (1975) brought back into fashion all the old works on these strange objects. He used them in a lot of different disciplines such as in physics (turbulence), geography ("How long

---

published on the original version in 1895 (Weierstrass, 1895).

[42] This word is derived from the Latin word *fractus* meaning *irregular, broken.*

is the Coast of Britain?"), astronomy (craters of the moon), biology, etc. , in order to show that objects without a tangent at any point are nothing but the rule in nature. The property being common to the different objects he treated, and thus the one implying a certain unity in nature, is the *self-similarity.* A fractal object is such that an enlarged part is similar to the whole object, which implies a scale invariance[43].

6 Irreversibility and microscopic chaos

In the context of statistical mechanics establishing relationships between the microscopic and macroscopic scales, we deal with microscopic chaos implied by the defocusing character of successive collisions between atoms and molecules. Let us consider the Brownian motion of a colloidal particle. The high-dimensional microscopic chaos of the surrounding fluid, characterized by the spectrum of Lyapunov exponents, induces a dynamical randomness given by the KS entropy, which is calculated by Pesin's identity (20). This huge dynamical randomness appears to be at the origin of the erratic motion of the Brownian particle. Consequently it gives us a new interpretation[44] of the observed stochastic processes in terms of a high-dimensional chaos in the microscopic Newtonian dynamics. Therefore the statistical mechanics of irreversible processes no longer needs stochastic models in order to describe the macroscopic irreversibility. The microscopic chaos provides deterministic systems which present a stochastic-like behavior without need of any stochastic assumption. Moreover, chaos and the sensitivity to initial conditions implying unpredictability for long times justifies the use of statistical mechanics, even for systems with a low-dimensional phase space.

In this perspective, during the last two decades, different theories have been developed establishing connections between irreversible phenomena (transport and chemical-reaction processes) and the chaotic properties of the microscopic dynamics. Among them we find the *thermostated-system approach* developed in the eighties by Nosé (1984a, 1984b), Hoover (1985, 1991), Evans and Morriss (Evans et al., 1990; Evans and Morriss, 1990), and others.

In this approach, nonequilibrium systems are defined as systems composed of particles submitted to interparticle forces, to external forces, but also to a fictitious nonHamiltonian force modeling the coupling to some hypothetical thermostat (Evans and Morriss, 1990). For instance, in order to study viscosity, the idea is to reproduce a Couette flow induced by a shearing force (see Fig. 3). Hence a velocity gradient is established in the system. However, this gradient leads to considerable viscous heating of the fluid and the energy of the system does not remain constant. To deal with this problem, it is necessary to introduce an internal thermostat – a fictitious frictional force – in order to keep constant the energy. Formally, for a 3-D system, in addition to the shearing force, we have to introduce a term with a thermostating multiplier α (Evans and Morriss, 1990)

---

[43] Several fundamental papers on the development of theory of fractals are reprinted in Elgar (1993).

[44] An experimental work has been done exhibiting the chaotic character of the microscopic dynamics by Gaspard et al. (1998).



Figure 3: Illustration of the shearing of the system inducing a velocity gradient $\frac{dv_x}{dy}$.

$$\begin{aligned}\frac{d\mathbf{q}_i}{dt} &= \frac{\mathbf{p}_i}{m} + \gamma y_i \\ \frac{d\mathbf{p}_i}{dt} &= \mathbf{F}_i - \gamma p_{yi} - \alpha \mathbf{p}_i \ , \end{aligned} \tag{21}$$

where γ is the vector $(\frac{dv_x}{dy}, 0)$, and $\mathbf{v}$ is the mean velocity. However, such dynamical systems violate Liouville theorem which asserts that the phase-space volumes are presented by the microscopic dynamics. This violation leads to fundamental problems for defining an entropy in nonequilibrium steady states. It thus appears as an artefact of a nonHamiltonian force that the phase-space volume visited by the system decreases in time and is expressed by the non-zero sum of Lyapunov exponents $\sum_{i=1}^{6N} \lambda_i < 0$. This phase-space contraction is introduced in the system through the presence of the thermostat (and the quantity α) which takes away the energy given to the system by shearing γ. It can be shown that the relation between the viscosity and the maximum and minimum Lyapunov exponents becomes for large systems (Evans et al., 1990)

$$\eta(N,\gamma) = \frac{-3nk_B T}{\gamma^2}\left[\lambda_{\max}(\gamma) + \lambda_{\min}(\gamma)\right] \ , \tag{22}$$

the N dependence of η and λ disappearing. The shear viscosity occurring in the Navier-Stokes equation is given by $\eta = \lim_{\gamma\to 0}\eta(\gamma)$. This expression relates the Lyapunov exponents to the viscosity coefficient because of the violation of Liouville's theorem by the artifitial nonHamiltonian systems. Therefore, this method cannot be used for Hamiltonian systems. Several works have shown that other ways exist to maintain a system out of equilibrium for instance by stochastic boundary conditions or by deterministic scattering (Klages et al., 2000) in which cases the relation (22) do not apply. Moreover, violating Louville's theorem creates problems in defining the entropy for nonequilibrium steady states.

Another theory developed in the nineties is the so-called *escape-rate formalism* (Gaspard, 1998; Dorfman, 1999) proposed in 1990 by Gaspard and Nicolis (1990) for diffusion. As for the previous method, the escape-rate formalism introduces nonequilibrium conditions. However, instead of modifying the Hamiltonian equations, we open the system in order to generate an escape process. More precisely, we impose absorbing boundary conditions at the statistical level of description, keeping the Hamiltonian character of the equations of motion themselves. The so-called *escape rate* is related to the studied transport coefficient on the one hand, and to the chaotic quantities of the microscopic dynamics on the other hand.

Let us illustrate the escape process by taking a simple example: the diffusion of a particle in the *periodic Lorentz gas* (see Fig. 4 (a)). This system consists of a particle of nonvanishing mass diffusing thanks to the successive elastic collisions with a set of two-dimensional fixed hard disks forming a hexagonal lattice. The escape process is introduced by limiting the scatterers in a direction, e.g. in the x-direction. It is said that the particle escapes when $|x| > \frac{L}{2}$ where L is an arbitrary parameter. The number of particles lying between the boundaries exponentially decreases in time at a rate given by the escape rate (see Figs. 4).

In 1995, Dorfman and Gaspard extented the escape-rate formalism to the other transport processes (Dorfman and Gaspard, 1995; Gaspard and Dorfman, 1995). Instead of imposing an external constraint like a shearing, as for the previous method, the nonequilibrium conditions are introduced by imposing



Figure 4: Lorentz gas with absorbing boundary conditions separated by the distance L. (a) Escape process of some particles after diffusion in the scatterer. (b) Exponential decrease in time of the number of particles lying into the limits defined by $-\frac{L}{2} \leq x \leq \frac{L}{2}$.

absorbing conditions

$$-\frac{\chi}{2} \leq G_{xy}(t) \leq +\frac{\chi}{2} \tag{23}$$

characterized by an arbitrary parameter χ, over the evolution the *Helfand moment* $G_{xy}(t)$ written in Eq. (19). When the Helfand moment reaches these absorbing boundaries, the system is said to be escaped. In consequence, considering a statistical ensemble of systems, an escape process appears, which is characterized by the escape rate. Since the system is in nonequilibrium, the support of the invariant measure is no longer the whole phase space, but a *fractal repeller*[45]. This fractal is constructed with the trajectories for which the Helfand moment remain forever into the absorbing boundaries (23). The escape rate plays the role of intermediate between the viscosity as a transport coefficient, and quantities of the microscopic chaos. Indeed, on the one hand, the viscosity can be written in terms of a *phenomenological* escape rate by considering the macroscopic escape process. On the other hand, the theory of dynamical systems gives a relationship between the escape rate and quantities such as the Lyapunov exponents and fractal dimensions, typical quantities of chaos (Ruelle, 1978; Eckmann and Ruelle, 1985; Beck and Schlögl, 1993). By identifying both escape rates, a relationship between the viscosity and the microscopic chaos is established and is written as (Dorfman and Gaspard, 1995; Gaspard and Dorfman, 1995)

---

[45] The fractal object is a repeller because of the instability of the dynamics due to its chaotic character.

$$\eta = \lim_{\chi \to \infty} \left(\frac{\chi}{\pi}\right)^2 \sum_{\lambda_i > 0} (1 - d_i) \ \lambda_i \ , \tag{24}$$

with the Lyapunov exponents λ_i and the associated partial dimensions d_i of the fractal repeller. Hence we can show how the microscopic chaos, responsible of the fractal character of the support of the invariant measure, controls the viscosity (Viscardy and Gaspard, 2003), or also the diffusion (Gaspard and Baras, 1995) as well as the reaction-diffusion processes (Claus and Gaspard, 2001; Claus et al., 2004).

The third approach is the *hydrodynamic-mode method* (Gaspard, 1998) developed during the nineties by Gaspard and coworkers (Gaspard, 1993; Gaspard, 1996). As discussed in Section 3, hydrodynamics describes the macroscopic dynamics of fluids in terms of equations governing the evolution of mass density, fluid velocity, and temperature, such as the Navier-Stokes equations (4) and the diffusion equation

$$\frac{\partial n}{\partial t} = D \nabla^2 n \ . \tag{25}$$

Thanks to the kinetic equation developed by Boltzmann nonequilibrium statistical mechanics is able to derive these phenomenological equations, while the Boltzmann equation is itself derived from Liouvillian dynamics, using the *Stosszahlantsatz*. The solutions of Eq.(25) called *hydrodynamic modes* are of the form

$$n_{\mathbf{k}}(\mathbf{r}, t) = \exp(s_{\mathbf{k}} t) \exp(i \mathbf{k} \cdot \mathbf{r}) \ , \tag{26}$$

each mode being characterized by a wavenumber $\mathbf{k}$. The hydrodynamic modes are spatially periodic of wavelength $\lambda = 2\pi/k$ with $k = ||\mathbf{k}||$. They decay exponentially in time because the corresponding eigenvalues are real and negative

$$s_{\mathbf{k}} = -Dk^2 \ . \tag{27}$$

The paradox between the reversible dynamics at the molecular level and the irreversible macroscopic phenomena that was exposed in section 2 can be studied in this context. Indeed the exponential decay of the modes (26) toward the thermodynamic equilibrium seems apparently incompatible with the Hamiltonian dynamics governing the motion of atoms and molecules, which is reversible and preserves phase-space volumes. Therefore, until recently the hydrodynamic modes were not described in terms of the Liouvillian dynamics, but at the intermediate level of the kinetic equations. Here, we briefly present a recent method based on the construction of hydrodynamic modes of diffusion in terms of the microscopic deterministic dynamics.

Let us consider the deterministic Liouvillian equation for Hamiltonian systems

$$\frac{\partial f}{\partial t} = \{H, f\} = \hat{L} f \ , \tag{28}$$

and the solution of this equation is

$$f_t(\mathbf{X}) = \hat{P}^t f_0(\mathbf{X}) \tag{29}$$

where $\hat{P}^t = \exp(\hat{L}t)$ is the so-called *Frobenius-Perron operator.*

Boundary conditions are required to solve the Liouville equation. This method does not use the simple periodic boundary conditions. Although we consider here N-particle systems periodically extended in position space and forming a lattice $\mathcal{L}^{\mathrm{N}}$, such as the Lorentz gas, the probability density is allowed to extend nonperiodically over the whole lattice so that the periodic boundary conditions does not apply and have to be replaced by the so-called *quasiperiodic boundary conditions*[46]. A Fourier transform must be carried out in position space to reduce the dynamics to the cell at the origin of the lattice ($\mathbf{l} = 0$). A wavenumber $\mathbf{k}$ is introduced which varies continuously in a Brillouin zone reciprocal to the lattice

$$f_{t,\mathbf{k}}(\mathbf{r}, \mathbf{p}) = \sum_{\mathbf{l} \in \mathcal{L}^{\mathrm{N}}} \exp(-i\mathbf{k} \cdot \mathbf{l}) f_t(\mathbf{r} + \mathbf{l}, \mathbf{p}) \ . \tag{30}$$

In particular the hydrodynamic mode of wavenumber $\mathbf{k}$ is an eigenstate of the operator $\hat{T}_{\mathbf{l}}$ of translation by the lattice vector $\mathbf{l}$

$$\hat{T}_{\mathbf{l}} \Psi_{\mathbf{k}} = \exp(i\mathbf{k} \cdot \mathbf{l}) \Psi_{\mathbf{k}} \ . \tag{31}$$

The wavenumber $\mathbf{k}$ characterizes the spatial periodicity of the observables and of the probability densities. Each Fourier component of the probability density evolves differently in time, which requires the introduction of a new Frobenius-Perron operator $\hat{R}_{\mathbf{k}}$ depending explicitly on the wavenumber $\mathbf{k}$. As the operator of translation $\hat{T}_{\mathbf{l}}$ commutes with the Frobenius-Perron operator $R_{\mathbf{k}}$ we can find an eigenstate common to both the spatial translations and the time evolution

---

[46] If the time evolution of the coordinates $\mathbf{\Gamma} = (\mathbf{r}, \mathbf{p}) = (\mathbf{r_1}, ..., \mathbf{r}_N, \mathbf{p_1}, ..., \mathbf{p}_N)$ of the N particles is governed by a first-order equations $\dot{\mathbf{\Gamma}} = \mathbf{F}(\mathbf{\Gamma})$, the vector field $\mathbf{F}$ is therefore symmetric under discrete position translations: $\mathbf{F}(\mathbf{r}, \mathbf{p}) = \mathbf{F}(\mathbf{r} + \mathbf{l}, \mathbf{p})$ with $\mathbf{l} \in \mathcal{L}^{\mathrm{N}}$.

$$\hat{R}_{\mathbf{k}}\Psi_{\mathbf{k}} = \exp(s_{\mathbf{k}}t)\Psi_{\mathbf{k}} \ . \tag{32}$$

At vanishing wavenumber, we recover the dynamics with periodic boundary conditions which admits an invariant probability measure describing the microcanonical equilibrium state. In contrast, an invariant probability measure no longer exists as soon as the wavenumber is non-vanishing. Instead, we find a complex measure which decays at a rate given by the so-called *Pollicott-Ruelle resonance* $s_{\mathbf{k}}$ (Gaspard, 1998). This measure defines the hydrodynamic mode of wavenumber $\mathbf{k}$ and the associated Pollicott-Ruelle resonance $s_{\mathbf{k}}$ gives the dispersion relation of the hydrodynamic mode (27).

Such microscopic hydrodynamic modes present an important difference with the phenomenological hydrodynamics. This difference holds in the fact that they are mathematical distributions or singular measures. The impossibility of constructing eigenstates in terms of functions has its origin in the pointlike character of the deterministic dynamics and in the property of dynamical instability. Therefore, in order to have a representation of the eigenstates we have to consider its cumulative function

$$F_{\mathbf{k}}(\xi) = \int_0^{\xi} \Psi_{\mathbf{k}}(\mathbf{\Gamma}_{\xi'})\, d\xi' \tag{33}$$

where $\mathbf{\Gamma}_{\xi'}$ is a curve of parameter ξ in the phase space.

The singular property of the eigenstates plays a fundamental role in the understanding of irreversible processes. Indeed it leads to the result that the cumulative function $F_{\mathbf{k}}(\xi)$ is fractal. In the case of diffusion in the periodic Lorentz gas, thanks to the thermodynamic formalism, it has been shown that the diffusion coefficient D is related to the Hausdorff dimension d_H of the cumulative function $F_{\mathbf{k}}(\xi)$ (Gaspard et al., 2001) as

$$D = \lambda \lim_{\mathbf{k}\to 0} \frac{d_H(\mathbf{k}) - 1}{\mathbf{k}^2} \tag{34}$$

where λ is the positive Lyapunov exponent. Let us mention that a similar study has been done in reactive-diffusion systems (Claus and Gaspard, 2004).

Such processes like diffusion and viscosity are typical irreversible phenomena. This irreversibility can be expressed by the production of entropy. It has been shown that the fractal structure of the diffusive modes plays a crucial role in the positivity of the entropy production in Hamiltonian systems (Gaspard, 1997; Gilbert et al., 2000). Indeed when the distribution functions are smooth, there is no change in the Gibbs entropy and no positive irreversible entropy production. The presence of the singular character therefore appears to be the fundamental element for an understanding of the second law of thermodynamics in terms of fractals.

7 Conclusions

In this paper, we have outlined the history of the origin of nonequilibrium statistical mechanics, treating the paradox between the reversible character of the microscopic dynamics and the irreversible transport processes, such as the viscosity. After having reviewed the development of hydrodynamics and the concept of viscosity, we have shown the development of the kinetic theory of gases mainly since Clausius and Maxwell who aimed to describe macroscopic phenomena in terms of the microscopic dynamics, and how appeared the first interpretation of the origin of irreversibility given by Boltzmann. Boltzmann's arguments were based on the probability that a system be in a certain microscopic state. For instance, consider an ideal gas in a box. Since the state in which all the particles are distributed in the whole vessel is infinitely more probable than the one in which the particles are concentrated only in the half part, it therefore seems natural that the system must tend to the homogeneous state and that we cannot observe the reverse case. Consequently, this interpretation is mainly *probabilist*.

We have presented an alternative description which has been developed in the last few decades. The rise of the theory of chaos and its application in dynamical system theory provide a *dynamical* intepretation of irreversibility in terms of the underlying microscopic chaos. Indeed, the typical instability of the dynamical chaos induces randomness which is a main characteristic of the transport processes (think of the erratic movement of a Brownian particle in a fluid). The successful results accumulated since the eighties confirm that microscopic chaos is an essential element at the origin of the irreversibility.

Acknowledgments. The author thanks Professors P. Gaspard and G. Nicolis for support and encouragement in this research, and the Fund for the Education to the Research in Industry and Agriculture (FRIA) for financial support.

References

Adams, H. (2004). *The Degradation of the Democratic Dogma.* Kessinger Publishing, Whitefish.

Balescu, R. (1960). Irreversible processes of ionized gases. *Phys. Fluids*, **3**:52–63.

Beck, C. and Schlögl, F. (1993). *Thermodynamics of Chaotic Systems: an Introduction.* Cambridge University Press, Cambridge.

Bernoulli, D. (1738). *Hydrodynamica, sive de viribus et motibus fluidorum commentarii.* Strasbourg.

Birkhoff, G. D. (1927). Dynamical systems. volume 9 of *Amer. Math. Soc. Colloquium Publ.*, American Mathematical Society, New York.

Blackwell, R. J. (1966). Descartes' laws of motion. *Isis*, **57**:220–234.

Blair, G. W. S. (1944). *A Survey of General and Applied Rheology.* Pitman Publishing Corp., New York.

Bogoliubov, N. N. (1946). Expansions into a series of powers of a small parameter in the theory of statistical equilibrium; kinetic equations. *J. Phys. USSR*, **10**:257, 265.

Boltzmann, L. (1868). Studien über das Gleichgewicht der lebendige Kraft zwischen bewegten materiellen Punkten. *Sitzungsberichte Akad. Wiss. Wien, Mathematisch-Naturwissenschafltliche Klasse*, **58**:517–560.

Boltzmann, L. (1872). Weitere Studien über das Wärmegleichgewicht unter Gasmolekülen. *Sitzungsberichte Akad. Wiss. Wien, Mathematisch-Naturwissenschafltliche Klasse*, **66**:275–370.

Boltzmann, L. (1884). Über die Eigenschaften monocylischer und anderer damit verwandter Systeme. *Sitzungsberichte, K. Akademie der Wissenschaften Wien, Mathematisch-Naturwissenschaftliche Klasse*, **90**:231–245.

Boltzmann, L. (1995). *Lectures on Gas Theory.* Dover, New York.

Born, M. and Green, H. S. (1946). A general kinetic theory of liquids. I. The molecular distribution functions. *Proc. Roy. Soc. lond. A*, **188**:10.

Boyle, R. (1660). *New experiments Physico-Mechanical, touching the Spring of the Air, and its Effects.* Oxford.

Brout, R. and Prigogine, I. (1956). Statistical mechanics of irreversible processes. Part VIII. General theory of weakly coupled systems. *Physica*, **22**:621–636.

Brush, S. G. (1957a). The development of the kinetic theory of gases. I. Herapath. *Annals of Sciences*, **13**:188–198.

Brush, S. G. (1957b). The development of the kinetic theory of gases. II. Waterston. *Annals of Sciences*, **13**:273–282.

Brush, S. G. (1962a). Development of the kinetic theory of gases. VI. Viscosity. *Am. J. of Phys.*, **30**:269–281.

Brush, S. G. (1962b). Theories of liquid viscosity. *Chem. Rev.*, **62**:513–548.

Brush, S. G. (1967). Thermodynamics and history: Science and culture in the nineteenth century. *The Graduate Journal*, **7**:477–565.

Brush, S. G. (1970). Interatomic forces and gas theory from Newton to Lennard-Jones. *Arch. Ration. Mech. Anal.*, **39**:1–29.

Brush, S. G. (1972). *Kinetic Theory, Volume 3: The Chapman-Enskog solution of the transport equation for moderately dense gases.* Pergamon Press, Oxford.

Brush, S. G. (1976). *The kind of motion we call heat: Book2. Statistical Physics and Irreversible processes.* North Holland, Amsterdam, New York & Oxford.

Brush, S. G. (1983). *Statistical Physics and the Atomic Theory of Matter, from Boyle and Newton to Landau and Onsager.* Princeton University Press, Princeton.

Brush, S. G. (1994). Geophysics. In Grattan-Guinness, I., editor, *Companion Encyclopedia of the History and Philosophy of the Mathematical Sciences*, pages 1183–1188. Routledge, London.

Brush, S. G. (1996). The reception of Mendeleev's periodic law in America and Britain. *Isis*, **87**:595–628.

Burich, K. R. (1987). Henry Adams, the second law of thermodynamics, and the course of history. *J. Hist. Ideas*, **48**:467–482.

Carnot, S. (1824). *Réflexions sur la puissance motrice du feu et sur les machines propres à développer cette puissance.* Bachelier, Paris. Translated in *Reflections on the motive power of fire by Sadi Carnot and other papers on the second law of thermodynamics by E. Clapeyron and R. Clausius*, E. Mendoza, Editor. Dover, New York, 1960.

Chabert, J.-L. (1994). The early history of fractals, 1870-1920. In Grattan-Guinness, I., editor, *Companion Encyclopedia of the History and Philosophy of the Mathematical Sciences*, pages 367–374. Routledge, London.

Chapman, S. (1917). The kinetic theory of simple and composite monoatomic gases: Viscosity, thermal conduction and diffusion. *Proc. of the Roy. Soc. of London A*, **93**:1–20. Reprinted in (Brush, 1972).

Chapman, S. and Cowling, T. G. (1970). *The Mathematical Theory of Non-uniform Gases.* Cambridge University Press, Cambridge.

Claus, I. and Gaspard, P. (2001). Fractals and dynamical chaos in a two-dimensional Lorentz gas with sinks. *Phys. Rev. E*, **63**:036227.

Claus, I. and Gaspard, P. (2004). The fractality of the relaxation modes in reaction-diffusion systems. *Physica D*, **187**:146–164.

Claus, I., Gaspard, P., and van Beijeren, H. (2004). Fractals and dynamical chaos in a random 2D Lorentz gas with sinks. *Physica D*, **187**:146–164.

Clausius, R. (1854). Über die mittlere Länge der Wege, welche bei der Molecularbewegung gasförmingen Körper von den einzelnen Molecülenzuruckgelegt werden; nebst einigen anderen Bemerkungen über die mechanische Wärmetheorie. *Ann. der Phys.*, **93**:481–506.

Clausius, R. (1856). On a modified form of the second fundamental theorem in the mechanical theory of heat. *Phil. Mag.*, **4**:81–98.

Clausius, R. (1858). Über die mittlere Länge der Wege, welche bei Molecularbewegung gasförming Körper von den einzelnen Molecülen zurückgelegt werden, nebst einigen anderen Bemerkungen über die mechanischen Wärmetheorie. *Ann. der Phys.*, **105**:239–258.

Clausius, R. (1865). Über verschiedene für die Anwendung bequeme Formen der Hauptgleichung der mechanischen. *Ann. der Phys.*, **125**:325–400.

Clausius, R. (1870). Über einen auf die Wärme anwendbaren mechanischen Satz. *Sitzungsberichte der Niedderrheinischen Gesellschaft, Bonn*, pages 114–119. Translated and reprinted in (**?**).

de Buffon, C. (1774). *Introduction à l'Histoire des Minéraux.* Paris.

de Saint-Venant, J. C. B. (1843). Note à joindre au Mémoire sur la dynamique des fluides, présenté le 14 avril 1834. *Comptes Rendus des Scéances de l'Académie des Sciences*, **17**:1240–1243.

Dorfman, J. R. (1999). *An Introduction to Chaos in Nonequilibrium Statistical Mechanics.* Cambridge University Press, Cambridge.

Dorfman, J. R. and Cohen, E. D. G. (1967). Difficulties in the kinetic theory of dense gases. *J. Math. Phys.*, **8**:282–297.

Dorfman, J. R. and Gaspard, P. (1995). Chaotic scattering theory of transport and reaction-rate coefficients. *Phys. Rev. E*, **51**:28–35.

Duhem, P. (1906). *La Théorie Physique, Son Objet, Sa Structure.* Chevalier et Rivière, Paris. Can be dowloaded from `http://www.ac-nancy-metz.fr/enseign/philo/textesph/`.

Eckmann, J. P. and Ruelle, D. (1985). Ergodic theory of chaos and strange attractors. *Rev. Mod. Phys.*, **57**:617–656.

Ehrenfest, P. and Ehrenfest, T. (1990). *The Conceptual Foundations of the Statistical Approach in Mechanics.* Dover, New York.

Einstein, A. (1905). Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssgkeiten suspendierten Teilchen. *Ann. d. Phys.*, **17**:549–560. Translated and reprinted in *Investigations on the theory of the brownian movement.* Dover, New York, 1956.

Elgar, G. A. (1993). *Classics on Fractals.* Addison-Wesley, MA.

Enskog, D. (1917). Kinetische Theorie der Vorgänge in mässig verdünnten Gasen. *published by Almqvist & Wiksells Boktryckeri-A.B., Uppsala.* Reprinted in (Brush, 1972).

Enskog, D. (1922). Kinetische Theorie der Wärmeleitung, Reibung und Selbstdiffusion in gewissen verdichtetenn Gasen und Flüssigkeiten. *Kungl. Sv. Vet. Akademiens Handlingar*, **63**(4). Reprinted in (Brush, 1972).

Euler, L. (1755). *Principes généraux de l'état d'équilibre des fluides; Principes généraux du mouvement des fluides; Continuation des recherches sur la théorie du mouvement des fluides.* Histoire de l'Académie de Berlin.

Evans, D. J., Cohen, E. D. G., and Morriss, G. P. (1990). Viscosity of a simple fluid from its maximal Lyapunov exponents. *Phys. Rev. A*, **42**:5990–5997.

Evans, D. J. and Morriss, G. P. (1990). *Statistical Mechanics of nonequilibrium Liquids*. Academic, London.

Feigenbaum, M. (1978). Quantitative universality for a class of nonlinear transformations. *J. Stat. Phys.*, **19**:25.

Fourier, J. (1819). Mémoire sur le Refroidissement Séculaire du Globe Terrestre. *Ann. Chimie*, **13**:418–437.

Freeman, D. (1974). The evolution theories of Charles Darwin and Herbert Spencer. *Current Anthropology*, **15**:211–237.

Garnett, C. (1882). *The Life of James Clerk Maxwell, with a selection from his correspondence and occasional writings, and a sketch of his contributions to science*. Macmillan.

Gaspard, P. (1993). What is the role of chaotic scattering in irreversible processes? *Chaos*, **3**:427–442.

Gaspard, P. (1996). Hydrodynamic modes as singular eigenstates of the Liouvillian dynamics: Deterministic diffusion. *Phys. Rev. E*, **53**:4379–4401.

Gaspard, P. (1997). Entropy production in open volume-preserving systems. *J. Stat. Phys.*, **88**:1215–1240.

Gaspard, P. (1998). *Chaos, Scattering and Statistical Mechanics*. Cambridge University Press, Cambridge.

Gaspard, P. and Baras, F. (1995). Chaotic scattering and diffusion in the Lorentz gas. *Phys. Rev. E*, **51**:5332–5352.

Gaspard, P., Briggs, M. E., Francis, M. K., Sengers, J. V., Gammon, R. W., Dorfman, J. R., and Calabrese, R. V. (1998). Experimental evidence for microscopic chaos. *Nature*, **394**:865–868.

Gaspard, P., Claus, I., Gibert, T., and Dorfman, J. (2001). Fractality of the hydrodynamic modes of diffusion. *Phys. Rev. Lett.*, **86**:1506–1509.

Gaspard, P. and Dorfman, J. R. (1995). Chaotic scattering theory, thermodynamic formalism, and transport coefficients. *Phys. Rev. E*, **52**:3525–3552.

Gaspard, P. and Nicolis, G. (1990). transport properties, lyapunov exponents, and entropy per unit time. *Phys. Rev. Lett.*, **65**:1693–1696.

Gibbs, J. (1902). *Elementary Principles in Statistical Mechanics.* Scribner, New York.

Gilbert, T., Dorfman, J., and Gaspard, P. (2000). Entropy production, fractals, and relaxation to equilibrium. *Phys. Rev. Lett.*, **85**:1606–1609.

Glansdorff, P. and Prigogine, I. (1971). *Thermodynamics of Structure, Stability, and Fluctuations.* Wiley-Interscience, New Yor.

Gleick, J. (1987). *Chaos: Making a new science.* Viking, New York.

Graham, T. (1846). On the motion of gases. *Phil. Trans. Roy. Soc. (London)*, **136**:573–631.

Graham, T. (1849). On the motion of gases. Part II. *Phil. Trans. Roy. Soc. (London)*, **139**:349–391.

Green, M. S. (1951). Brownian motion in a gas of noninteracting molecules. *J. Chem. Phys.*, **19**:1036–1046.

Green, M. S. (1960). Comment on a paper of Mori on time-correlation expressions for transport properties. *Phys. Rev.*, **119**:829–830.

Hadamard, J. (1898). Les surfaces à courbes opposées et leurs lignes géodésiques. *J. Math. Pur. Appl.*, **4**:27–73.

Hagenbach, E. (1860). Über die Bestimmung der Zähigkeit einer Flüssigkeit durch den Aufsluss aus Röhren. *Pogg. Ann.*, **109**:385–426.

Hanley, H. J. M. and Cohen, E. D. G. (1976). Analysis of the transport coefficients for simple dense fluids: The diffusion and bulk viscosity coefficients. *Physica A*, **83**:215–232.

Hanley, H. J. M., McCarty, R. D., and Cohen, E. D. G. (1972). Analysis of the transport coefficients for simple dense fluids: Application of the modified enskog theory. *Physica*, **60**:322–356.

Hausdorff, F. (1919). Dimension und äußeres Maß. *Mathematische Annalen*, **79**:157–179. Translated and reprinted in (Elgar, 1993).

Hirschfelder, J. O., Curtiss, C. F., and Bird, R. B. (1954). *Molecular Theory of Gases and Liquids*. John Wiley & Sons, New York.

Holton, G. and Brush, S. G. (1985). *Introduction to concepts and theories in physical science*. Princeton University Press, Princeton.

Hoover, W. G. (1985). Canonical dynamics: Equilibrium phase-space distributions. *Phys. Rev. A*, **31**:1695–1697.

Hoover, W. G. (1991). *Computational Statistical Mechanics*. Elsevier, New York.

Hove, L. V. (1955). Quantum-mechanical perturbations giving rise to a statistical transport equation. *Physica*, **21**:517–540.

Hove, L. V. (1957). The approach to equilibrium in quantum statistics: A perturbation treatment to general order. *Physica*, **23**:441–480.

Hove, L. V. (1959). The ergodic behaviour of quantum many-body systems. *Physica*, **25**:268–276.

Hunt, B. R. and Yorke, J. A. (1993). Maxwell on chaos. *Nonlinear Science Today*, 3:2–4.

Jeans, J. (1903). *Phil. Mag.*, **4**:585.

Jeans, J. (1904). *The Dynamical Theory of Gases*. Cambridge University Press, Cambridge.

Joule, J. (1847). On matter, living force, and heat. *Manchester Courier*. Reprinted in (**?**).

Kamgar-Parsi, B. and Sengers, J. V. (1983). Logarithmic density dependence of the transport properties of gases. *Phys. Rev. Lett.*, **51**:2163–2166.

Karim, S. M. and Rosenhead, L. (1952). The second coefficient of viscosity of liquids and gases. *Rev. Mod. Phys.*, **24**:108–116.

Kestin, J., Korfali, O., and Sengers, J. V. (1980). Density expansion of the viscosity of carbon dioxide near the critical temperature. *Physica A*, **100**:335–348.

Kestin, J., Korfali, O., and Sengers, J. V. (1981). Density dependence and composition dependence of the viscosity of neon-helium and neon-argon mixtures. *Physica A*, **106**:415–442.

Kestin, J., Paykoç, E., and Sengers, J. V. (1971). On the density expansion for viscosity in gases. *Physica*, **54**:1–19.

Kirkwood, J. G. (1946). The statistical mechanical theory of transport processes. I. General theory. *J. Chem. Phys.*, **14**:180–201.

Klages, R., Rateitschak, K., and Nicolis, G. (2000). Thermostating by deterministic scattering: Construction of nonequilibrium steady states. *Phys. Rev. Lett.*, **84**:4268–4271.

Koch, H. (1904). On a continuous curve without tangent construtible from elementary geometry. *Arkiv för Matematik, Astronomi och Fysik*, **1**:678–704.

Koyré, A. (1957). *From the Closed World to the Infinite Universe.* John Hopkins Press, Baltimore.

Krönig, A. (1856). Grundzuge einer Theorie der Gase. *Ann. d. Phys.*, **99**:315–322.

Krylov, N. S. (1979). *Works on the Foundations of Statistical Physics.* Princeton University Press, Princeton.

Kubbinga, H. (1990). Thermodynamics, molecularism, and positivism. In K. Martinás, L. R. and Szegedi, P., editors, *Thermodynamics: History and Philosophy*, pages 404–415. World Scientific, Singapore.

Kubo, R. (1957). Statistical-mechanical theory of irreversible processes. I. General theory and simple applications to magnetic and conduction problems. *J. Phys. Soc. Jpn.*, **12**:570–586.

Landau, L. D. and Lifshitz, E. M. (1959). *Fluid Mechanics.* Pergamon Press, Oxford.

Leicester, H. M. (1956). *The historical background of chemistry.* Dover, New York.

Lenard, A. (1960). On Bogoliubov's kinetic equation for a spatially homogeneous plasma. *Ann. Phys.*, **3**:390.

Lennard-Jones, J. E. (1931). Cohesion. *Proc. Phys. Soc. London*, **43**:461–482.

Lorenz, E. (1963). Deterministic nonperiodic flow. *J. Atm. Sciences*, **20**:130–141.

Loschmidt, J. (1865). Zur Grösse der Luftmolecüle. *Sitzb. Akad. Wiss. Wien*, **395-413**:395–413. Translated and reprinted by Porterlied and Kruse (1995).

Lyapunov, A. (1892). General problems in the stability of motion (in russian). *Comm. Soc. Math. Kharkow*. Reprinted in *Ann. Math. Studies* **17**, Princeton (1949).

Madigosky, W. M. (1967). Density dependence of the bulk viscosity in argon. *J. Chem. Phys.*, **46**:4441–4444.

Mandelbrot, B. (1975). *Les Objets fractals: Forme, hasard et dimension.* Flammarion, Paris.

Markham, J. J., Beyer, R. T., and Lindsay, R. B. (1951). Absorption of sound in fluids. *Rev. Mod. Phys.*, **23**:353–411.

Maxwell, J. (1860). Illustrations of the dynamical theory of gases. *Phil. Mag.*, **19**:19–32.

Maxwell, J. (1866). On the viscosity or internal friction of air and other gases. *Phil. Trans.*, **156**:249–268.

Mayer, R. (1842). Bemerkungen über die Kräfte der unbelebten Natur. *Annalen der Chemie und Pharmacie*, **42**:233–240. Translated and reprinted in (**?**).

Mendeleev, D. (1869a). The relation between the properties and atomic weights of the elements. *Journal of the Russian Chemical Society*, **1**:60–77.

Mendeleev, D. (1869b). The relation between the properties and atomic weights of the elements (abstract of (Mendeleev, 1869a)). *Zeitscrift für Chemie*, **12**:405–406.

Michels, A. and Gibson, R. O. (1931). The measurement of the viscosity of gases at high pressures: The viscosity of nitrogen to 1000 atms. *Proc. Roy. Soc. London A*, **34**:288–307.

Mikhailov, G. K. (1994). Hydrodynamics and hydraulics. In Grattan-Guinness, I., editor, *Companion Encyclopedia of the History and Philosophy of the Mathematical Sciences*, pages 1006–1022. Routledge, London.

Mohr, P. J. and Taylor, B. N. (1998). CODATA recommended values of the fundamental physical constants: 1998. *Rev. Mod. Phys.*, **71**:351–495.

Mori, H. (1958). Statistical-mechanical theory of transport in fluids. *Phys. Rev.*, **112**:1829–1842.

Naugle, D. G. and Squire, C. F. (1965). Ultrasonic attenuation in liquid argon. *J. Chem. Phys.*, **42**:3725.

Navier, L. M. H. (1823). Mémoire sur les lois du mouvement des fluides. *Mémoires de l'Académie Royale des Sciences de l'Institut de France*, **6**:389–440.

Newton, I. (1952). *Opticks.* Dover, New York.

Newton, I. (1999). *The Principia : mathematical principles of natural philosophy.* University of California Press, Berkeley.

Nicolis, G. (1995). *Introduction to Nonlinear Science.* Cambridge University Press, Cambridge.

Nicolis, G. and Prigogine, I. (1977). *Self-Organization in Nonequilibrium Systems: From Dissipative Structures to Order through Fluctuations.* Wiley-Interscience, New York.

Nosé, S. (1984a). A molecular dynamics method for simulations in the canonical ensemble. *Mol. Phys.*, **52**:255–268.

Nosé, S. (1984b). A unified formulation of the constant temperature molecular dynamics methods. *J. Chem. Phys.*, **81**:511.

Ott, E. (1993). *Dynamical systems.* Cambridge University Press, Cambridge.

Pauli, W. (1928). Festschrift zum 60. In Hirzel, editor, *Geburstage A. Sommerfeld*, page 30. Leipzig.

Perrin, J. (1908). L'agitation moléculaire et le mouvement brownien. *Comptes Rendus*, **146**:967–970.

Perrin, J. (1991). *Les Atomes.* Flammarion, Paris. Translated in *The Atoms*, Ox Bow Press, Woodbridge, 1990.

Pesin, Y. B. (1977). Characteristic Lyapunov exponents and smooth ergodic theory. *Russ. Math. Surveys*, **32**:55–114.

Poincaré, H. (1892). *Les Méthodes Nouvelles de la Mécanique Céleste.* Gauthiers-Villard, Paris.

Poincaré, H. (1908). *Science et Méthode.* Flammarion, Paris.

Poiseuille, J. L. M. (1840a). Recherches expérimentales sur le mouvement des liquides dans les tubes de très petits diamètres. *Comptes Rendus*, **11**:961–967.

Poiseuille, J. L. M. (1840b). Recherches expérimentales sur le mouvement des liquides dans les tubes de très petits diamètres (suite). *Comptes Rendus*, **11**:1041–1048.

Poiseuille, J. L. M. (1846). Recherches expérimentales sur le mouvement des liquides dans les tubes de très petits diamètres. *Mémoire des Savants Etrangers*, **9**:463–544.

Poisson, S. (1831). Mémoire sur les équations générales de l'équilibre et du mouvement des corps élastiques et des fluides. *Journal de l'Ecole Polytechnique*, **13**:1–174.

Pullman, B. (1995). *L'atome dans l'histoire de la pensée humaine.* Fayard, Paris. Translated in *The atom in the history of human thought.* Oxford University Press, Oxford, 1998.

Rouse, H. and Ince, S. (1957). *History of Hydraulics.* Iowa Institute of Hydraulic Research, Iowa.

Rousseau, P. (1945). *Histoire de la Science.* Librairie Arthème Fayard, Paris.

Ruelle, D. (1978). *Thermodynamic Formalism.* Addison-Wesley Publishing Co., New York.

Russo, L. (2004). *The Forgotten Revolution.* Springer-Verlag, Berlin.

Sengers, J. V. (1965). Thermal conductivity and viscosity of simple fluids. *Int. J. Heat. Transfer*, **8**:1103–1116.

Sengers, J. V. (1966). Divergence in the density expansion of the transport coefficients of a two-dimensional gas. *Phys. Fluids*, **9**:1685–1696.

Sengers, J. V. (1968). Transport properties of compressed gases. In Eringen, A. C., editor, *Recent Advances in Engineering Science*, pages 153–196. Gordon and Breach, New York.

Sinai, Y. G. (1970). Dynamical systems with elastic reflections. ergodic properties of dispersing billiards. *Russian Math. Surveys*, **25**:137–189.

Smale, S. (1967). Differentiable dynamical systems. *Bull. Amer. Math. Soc.*, **73**:747.

Stokes, G. G. (1845). On the theories of the internal friction of fluids in motion, nd of the equilibrium and motion of elastic solids. *Trans. Cambridge Phil. Soc.*, **8**:287. Reprinted in G. G. Stokes, *Mathematical and Physical Papers: Vol.1*, p.75-129. Cambridge University Press, Cambridge, 1880.

Stokes, G. G. (1846). Report on recent researches in hydrodynamics. *Report of the British Association for 1846*, **Part I**:1. *Mathematical and Physical Papers: Vol.1*, p.157-187. Cambridge University Press, Cambridge, 1880.

Takagi, O. (1903). A simple example of the continuous function without derivative. *Proc. Phys. Math. Soc. Japan. Ser. II*, **1**:176.

Thompson, P. A. (1972). *Compressible-fluid Dynamics.* McGraw-Hill, New York.

Tisza, L. (1942). Supersonic absorption and Stokes' viscosity relation. *Phys.Rev.*, **61**:531–536.

Tokaty, G. A. (1994). *A History and Philosophy of Fluid Mechanics.* Dover, New York.

Uehling, E. A. and Uhlenbeck, G. E. (1933). Transport phenomena in Einstein-Bose and Fermi-Dirac gases. I. *Phys. Rev*, **43**:552–561.

van der Waals, J. (1873). *Over de continuiteit van den gas- en vloeisoftoestand.* Stijthoff, Leiden.

Viscardy, S. (2005). *Viscosity and Microscopic Chaos : The Helfand-moment Approah.* PhD thesis, Center for Nonlinear phenomena and Complex

Systems, Université Libre de Bruxelles, Brussels, Belgium. e-print: cond-mat/0510445.

Viscardy, S. and Gaspard, P. (2003). Viscosity in the escape-rate formalism. *Phys. Rev. E*, **68**:041205.

Warburg, E. and von Babo, L. (1882). *Ann. d. Physik*, **17**:390.

Weierstrass, K. (1895). Über continuirliche Functionen eines reellen Arguments, die für keinen Werth des Letzteren einen bestimmten Differentialquotienten besitzen. *Mathematische Werke (Königlich Preussischen Akademie der Wissenschaften, Berlin, 1895).* Reprinted by Elgar (1993).

Wiedemann, G. (1856). Über die Bewegung der Flüssigkeiten im Kreise der geschlossenen galvanischen Säule und ihre Beziehungen zur Elektrolyse. *Pogg. Ann.*, **99**:177–233.

Yvon, J. (1935). *La Théorie Statistique des Fluides et l'Equation d'Etat.* Hermann, Paris.